

Package ‘ALUES’

January 20, 2025

Type Package

Title Agricultural Land Use Evaluation System

Version 0.2.1

Date 2022-01-10

Maintainer Al-Ahmadgaid B. Asaad <alahmadgaid@gmail.com>

Author Al-Ahmadgaid B. Asaad [aut, cre]

(<<https://orcid.org/0000-0003-3784-8593>>),

Arnold R. Salvacion [aut] (<<https://orcid.org/0000-0001-8868-2226>>),

Bui Tan Yen [aut]

Description Evaluates land suitability for different crops production.

The package is based on the Food and Agriculture Organization (FAO) and the International Rice Research Institute (IRRI) methodology for land evaluation.

Development of ALUES is inspired by similar tool for land evaluation, Land Use Suitability Evaluation Tool (LUSET). The package uses fuzzy logic approach to evaluate land suitability of a particular area based on inputs such as rainfall, temperature, topography, and soil properties. The membership functions used for fuzzy modeling are the following: Triangular, Trapezoidal and Gaussian. The methods for computing the overall suitability of a particular area are also included, and these are the Minimum, Maximum and Average. Finally, ALUES is a highly optimized library with core algorithms written in C++.

License MIT + file LICENSE

LazyData true

LinkingTo Rcpp

Encoding UTF-8

Depends R (>= 3.5.0), Rcpp (>= 0.10.6)

Suggests testthat, markdown, knitr, microbenchmark, ggmap, raster, reshape2

RoxygenNote 7.1.1

VignetteBuilder knitr

URL <https://github.com/alstat/ALUES/>

BugReports <https://github.com/alstat/ALUES/issues/>

NeedsCompilation yes

Repository CRAN

Date/Publication 2022-01-12 21:12:44 UTC

Contents

ALUES-package	7
ALFALFASoil	7
ALFALFATemp	8
ALFALFATerrain	9
ALFALFAWater	9
AVOCADOSoil	10
AVOCADOTemp	11
AVOCADOTerrain	11
AVOCADOWater	12
BAMBOOSoil	12
BAMBOOTemp	13
BAMBOOTerrain	13
BAMBOOWater	14
BANANASoil	14
BANANATemp	15
BANANATerrain	16
BANANAWater	16
BARLEYSoil	17
BARLEYTemp	18
BARLEYTerrain	18
BARLEYWater	19
BEANCASoil	19
BEANCATemp	20
BEANCATerrain	21
BEANCAWater	21
BEANSSoil	22
BEANSTemp	23
BEANSTerrain	23
BEANSWater	24
CABBAGESoil	24
CABBAGETemp	25
CABBAGETerrain	26
CABBAGEWater	26
CARROTSSoil	27
CARROTSTemp	28
CARROTSTerrain	28
CARROTSWater	29
CASHEWSoil	29
CASHEWTemp	30
CASHEWTerrain	31
CASHEWWater	31

CASSAVA	Soil	32
CASSAVA	Temp	33
CASSAVA	Terrain	33
CASSAVA	Water	34
CHICKPEA	Soil	34
CHICKPEA	Temp	35
CHICKPEA	Terrain	36
CHICKPEA	Water	36
CINNAMON	Soil	37
CINNAMON	Temp	38
CINNAMON	Terrain	38
CINNAMON	Water	39
CITRUS	Soil	39
CITRUS	Temp	40
CITRUS	Terrain	41
CITRUS	Water	41
COCOA	Soil	42
COCOA	Temp	43
COCOA	Terrain	43
COCOA	Water	44
COCONUT	Soil	44
COCONUT	Temp	45
COCONUT	Terrain	46
COCONUT	Water	46
COFFEE	Soil	47
COFFEE	Temp	48
COFFEE	Terrain	48
COFFEE	Water	49
COFFEER	Soil	49
COFFEER	Temp	50
COFFEER	Terrain	51
COFFEER	Water	51
COTTON	Soil	52
COTTON	Temp	53
COTTON	Terrain	53
COTTON	Water	54
COWPEA	Soil	55
COWPEA	Temp	55
COWPEA	Terrain	56
COWPEA	Water	57
CUCUMBER	Soil	57
CUCUMBER	Temp	58
CUCUMBER	Terrain	59
CUCUMBER	Water	59
GROUNDNUT	Soil	60
GROUNDNUT	Temp	61
GROUNDNUT	Terrain	61
GROUNDNUT	Water	62

GUAVASoil	62
GUAVATemp	63
GUAVATerrain	64
GUAVAWater	64
LaoCaiLT	65
LaoCaiTemp	65
LaoCaiWater	66
LONGANSoil	67
LONGANTemp	68
LONGANTerrain	68
LONGANWater	69
MAIZESoil	69
MAIZETemp	70
MAIZETerrain	71
MAIZEWater	71
MANGOSoil	72
MANGOTemp	73
MANGOTerrain	73
MANGOWater	74
MarinduqueLT	74
MarinduqueTemp	75
MarinduqueWater	76
MILLETSSoil	76
MILLETSTemp	77
MILLETSTerrain	78
MILLETWater	78
OILPALMSoil	79
OILPALMTemp	80
OILPALMTerrain	80
OILPALMWater	81
OLIVESSoil	81
OLIVESTemp	82
OLIVESTerrain	83
OLIVESWater	83
ONIONSoil	84
ONIONTemp	85
ONIONTerrain	85
ONIONWater	86
overall_suit	86
PAPAYASoil	88
PAPAYATemp	88
PAPAYATerrain	89
PAPAYAWater	89
PEACHSoil	90
PEACHTemp	91
PEACHTerrain	91
PEACHWater	92
PEARSoil	92

PEARTemp	93
PEARTerrain	93
PEARWater	94
PEASoil	95
PEATemp	95
PEATerrain	96
PEAWater	97
PEPPERGRSoil	97
PEPPERGRTemp	98
PEPPERGRTerrain	98
PEPPERGRWater	99
PERSIMMONSoil	100
PERSIMMONTemp	101
PERSIMMONTerrain	101
PERSIMMONWater	102
PINEAPPLESoil	102
PINEAPPLETemp	103
PINEAPPLETerrain	103
PINEAPPLEWater	104
PLUMSoil	105
PLUMTemp	106
PLUMTerrain	106
PLUMWater	107
POTATOSoil	107
POTATOSWSoil	108
POTATOSWTemp	109
POTATOSWTerrain	109
POTATOSWWater	110
POTATOTemp	110
POTATOTerrain	111
POTATOWater	112
RICEBRSoil	112
RICEBRTemp	113
RICEBRTerrain	114
RICEBRWater	114
RICEIWSoil	115
RICEIWTemp	116
RICEIWTerrain	116
RICEIWWater	117
RICENFSoil	118
RICENFTemp	119
RICENFTerrain	119
RICENFWater	120
RICEURSoil	121
RICEURTemp	121
RICEURTerrain	122
RICEURWater	123
RUBBERSoil	123

RUBBERTemp	124
RUBBERTerrain	125
RUBBERWater	125
SAFFLOWERSoil	126
SAFFLOWERTemp	126
SAFFLOWERTerrain	127
SAFFLOWERWater	127
SESAMESoil	128
SESAMETemp	129
SESAMETerrain	129
SESAMEWater	130
SORGHUMSoil	130
SORGHUMTemp	131
SORGHUMTerrain	132
SORGHUMWater	132
SOYASoil	133
SOYATemp	134
SOYATerrain	134
SOYAWater	135
SUGARCANESoil	136
SUGARCANETemp	137
SUGARCANETerrain	137
SUGARCANEWater	138
suit	138
suitability	141
SUNFLOWERSoil	142
SUNFLOWERTemp	143
SUNFLOWERTerrain	144
SUNFLOWERWater	144
TEASoil	145
TEATemp	146
TEATerrain	146
TEAWater	147
TOBACCOSoil	148
TOBACCOTemp	149
TOBACCOTerrain	149
TOBACCOWater	150
TOMATOSoil	150
TOMATOTemp	151
TOMATOTerrain	152
TOMATOWater	152
WATERMELONSoil	153
WATERMELONTemp	154
WATERMELONTerrain	154
WATERMELONWater	155
WHEATSoil	155
WHEATTemp	156
WHEATTerrain	157

ALUES-package *Agricultural Land Use Evaluation System*

Description

Agricultural Land Use Evaluation System (ALUES) is a package that evaluates land suitability for different crop production. The package is based on the Food and Agriculture Organization (FAO) and the International Rice Research Institute (IRRI) methodology for land evaluation. Development of ALUES is inspired by similar tool for land evaluation, Land Use Suitability Evaluation Tool (LUSET). The package uses fuzzy logic approach to evaluate land suitability of a particular area based on inputs such as rainfall, temperature, topography, and soil properties. The membership functions used for fuzzy modeling are the following: Triangular, Trapezoidal, Gaussian, Sigmoidal and custom models with functions that can be defined by the user. The package also aims on complicated methods like considering more than one fuzzy membership function on different suitability class. The methods for computing the overall suitability of a particular area are also included, and these are the Minimum, Maximum, Product, Sum, Average, Exponential and Gamma. Finally, ALUES utilizes the power of Rcpp library for efficient computation.

Author(s)

Al-Ahmadgaid B. Asaad <alahmadgaid@gmail.com> (maintainer)

Arnold R. Salvacion <arsalvacion@gmail.com>

Bui Tan Yen

ALFALFASoil *Alfalfa soil requirement for land evaluation*

Description

A dataset containing the soil characteristics of the crop requirements for farming Alfalfa.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- SoilTe - 12 classes of soil texture (Soil Taxonomy)
- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

ALFALFATemp

Alfalfa temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Alfalfa.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

ALFALFATerrain *Alfalfa terrain requirement for land evaluation*

Description

A dataset containing the terrain characteristics of the crop requirements for farming Alfalfa.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

ALFALFAWater *Alfalfa water requirement for land evaluation*

Description

A dataset containing the water characteristics of the crop requirements for farming Alfalfa.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- CropLen - Length of growing period (days)
- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

AVOCADOSoil

Avocado soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Avocado.

Format

A data frame with 10 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pH_{H2O} - pH H₂O
- OC - Organic carbon (%)
- EC_{dS} - EC_e (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 AVOCADOTemp

Avocado temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Avocado.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMinXm - Avarage minimum temperature of coldest month (C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 AVOCADOTerrain

Avocado terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Avocado.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

AVOCADOWater

Avocado water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Avocado.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BAMBOOSoil

Bamboo soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Bamboo.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- SoilDpt - Soil depth (cm)
- OC - Organic carbon (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)
- pHH2O - pH H2O

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BAMBOOTemp

Bamboo temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Bamboo.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BAMBOOTerrain

Bamboo terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Bamboo.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BAMBOOWater

Bamboo water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Bamboo.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BANANASoil

Banana soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Banana.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)

- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BANANATemp

Banana temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Banana.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TyMaxAv - Mean annual maximum temperature (°C)
- TmMinXm - Avarage minimum temperature of coldest month (C)
- TmMinXmAb - Absolute min temp. coldest month (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BANANATerrain

Banana terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Banana.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BANANAWater

Banana water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Banana.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BARLEYSoil

Barley soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Barley.

Format

A data frame with 13 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 BARLEYTemp

Barley temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Barley.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmAv3 - Mean temp. of the flowering stage (°C)
- TmAv4 - Mean temp. of the ripening stage (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 BARLEYTerrain

Barley terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Barley.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BARLEYWater

Barley water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Barley.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BEANCASoil

Castor Beans soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Castor Beans.

Format

A data frame with 9 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BEANCATemp

Castor Beans temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Castor Beans.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv1 - Mean temp. of the initial stage(C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BEANCATerrain

Castor Beans terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Castor Beans.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BEANCAWater

Castor Beans water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Castor Beans.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmhAv3 - Relative humidity of maturation Stage (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BEANSSoil

Beans soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Beans.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BEANSTemp

Beans temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Beans.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BEANSTerrain

Beans terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Beans.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BEANSWater

Beans water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Beans.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv3 - Relative humidity of maturation Stage (%)
- WmnN2 - n/N develop. Stage (2nd month)
- WmnN4 - n/N maturation stage (4th month)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CABBAGESoil

Cabbage soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Cabbage.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CABBAGETemp

Cabbage temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Cabbage.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)
- TdDiff - Temp difference day/night (C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CABBAGETerrain *Cabbage terrain requirement for land evaluation*

Description

A dataset containing the terrain characteristics of the crop requirements for farming Cabbage.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CABBAGEWater *Cabbage water requirement for land evaluation*

Description

A dataset containing the water characteristics of the crop requirements for farming Cabbage.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CARROTSSoil

Carrots soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Carrots.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 CARROTSTemp

Carrots temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Carrots.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 CARROTSTerrain

Carrots terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Carrots.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CARROTSWater

Carrots water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Carrots.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CASHEWSoil

Cashew soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Cashew.

Format

A data frame with 8 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CASHEWTemp

Cashew temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Cashew.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TyMaxAv - Mean annual maximum temperature (°C)
- TmMinXm - Average minimum temperature of coldest month (C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CASHEWTerrain

Cashew terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Cashew.

Format

A data frame with 8 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CASHEWater

Cashew water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Cashew.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CASSAVASoil

Cassava soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Cassava.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm1 - Coarse fragment in surface (Vol.%)
- CFragm2 - Coarse fragment in depth (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CASSAVATemp

Cassava temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Cassava.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TyMaxAv - Mean annual maximum temperature (°C)
- TmMinXmAb - Absolute min temp. coldest month (°C)
- TgAv - Mean temperature of the growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CASSAVATerrain

Cassava terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Cassava.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 CASSAVAWater

Cassava water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Cassava.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WmnN5 - n/N of the 5 dryest months

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 CHICKPEASoil

Chickpea soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Chickpea.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CHICKPEATemp

Chickpea temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Chickpea.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CHICKPEATerrain

Chickpea terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Chickpea.

Format

A data frame with 8 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CHICKPEAWater

Chickpea water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Chickpea.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CINNAMONSoil

Cinnamon soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Cinnamon.

Format

A data frame with 9 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CINNAMONTemp	<i>Cinnamon temp requirement for land evaluation</i>
--------------	--

Description

A dataset containing the temp characteristics of the crop requirements for farming Cinnamon.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CINNAMONTerrain	<i>Cinnamon terrain requirement for land evaluation</i>
-----------------	---

Description

A dataset containing the terrain characteristics of the crop requirements for farming Cinnamon.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- SlopeD - Slope (degree, 6 classes)
- Flood - Flooding

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CINNAMONWater

Cinnamon water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Cinnamon.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CITRUSSoil

Citrus soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Citrus.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)

- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CITRUSTemp

Citrus temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Citrus.

Format

A data frame with 8 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TmMax38 - No of months with mean temp. > 38 °C
- TmMin13 - No of months with mean temp. < 13 °C
- TyMinAb - Absolute minimum temperature (°C)
- TyMinAb - Absolute minimum temperature (°C)
- TyMinAb - Absolute minimum temperature (°C)
- TyMinAb - Absolute minimum temperature (°C)
- TmAv3 - Mean temp. of the flowering stage (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CITRUSTerrain

Citrus terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Citrus.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CITRUSWater

Citrus water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Citrus.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WmhColdXm - Relative humidity of coldest month if frost (%)
- WmhAv4 - Relative humidity at harvest stage (%)
- WmhAv3 - Relative humidity of maturation Stage (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COCOASoil

Cocoa soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Cocoa.

Format

A data frame with 10 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyphs - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COCOATemp

Cocoa temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Cocoa.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TyMaxAv - Mean annual maximum temperature (°C)
- TyMaxAv - Mean annual maximum temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COCOATerrain

Cocoa terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Cocoa.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 COCOAWater

Cocoa water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Cocoa.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WmhDryXm - Mean rel. humidity dryest month (%)
- WmhDryXm - Mean rel. humidity dryest month (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 COCONUTSoil

Coconut soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Coconut.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- OC - Organic carbon (%)
- ECemh - ECe (mmhos/cm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COCONUTTemp

Coconut temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Coconut.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COCONUTTerrain *Coconut terrain requirement for land evaluation*

Description

A dataset containing the terrain characteristics of the crop requirements for farming Coconut.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COCONUTWater *Coconut water requirement for land evaluation*

Description

A dataset containing the water characteristics of the crop requirements for farming Coconut.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WyhAv - Mean annual rel. humidity (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COFFEEARSoil

Arabica Coffee soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Arabica Coffee.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 COFFEEARTemp

Arabica Coffee temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Arabica Coffee.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TyMaxAv - Mean annual maximum temperature (°C)
- TdMinXm - Mean daily minimum temperature of coldest month (°C)
- TyAv - Mean annual temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 COFFEEARTerrain

Arabica Coffee terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Arabica Coffee.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COFFEEARWater

Arabica Coffee water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Arabica Coffee.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WmhDryXm - Mean rel. humidity dryest month (%)
- WmnN5 - n/N of the 5 dryest months

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COFFEEROSoil

Robusta Coffee soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Robusta Coffee.

Format

A data frame with 10 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COFFEEROTemp

Robusta Coffee temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Robusta Coffee.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TyMaxAv - Mean annual maximum temperature (°C)
- TdMinXm - Mean daily minimum temperature of coldest month (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COFFEEROTerrain *Robusta Coffee terrain requirement for land evaluation*

Description

A dataset containing the terrain characteristics of the crop requirements for farming Robusta Coffee.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COFFEEROWater *Robusta Coffee water requirement for land evaluation*

Description

A dataset containing the water characteristics of the crop requirements for farming Robusta Coffee.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WmhDryXm - Mean rel. humidity driest month (%)
- WmnN5 - n/N of the 5 driest months

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COTTONSoil

Cotton soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Cotton.

Format

A data frame with 13 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragn - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECemh - ECe (mmhos/cm)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COTTONTemp

Cotton temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Cotton.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMaxAv - Mean max temp. of growing cycle (°C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TdAvg3 - Mean DAY temp. of flowering stage (°C)
- TdMinN3 - Mean Night temp. of flowering stage (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COTTONTerrain

Cotton terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Cotton.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- Slope - nan

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COTTONWater

Cotton water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Cotton.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv5 - Mean precipitation of fifth month (mm)
- WmAv6 - Precipitation of ripening stage (mm)(6th month)
- WmhAv3 - Relative humidity of maturation Stage (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COWPEASoil

Cowpea soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Cowpea.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COWPEATemp

Cowpea temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Cowpea.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)
- TyMinAv - Mean annual minimum temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COWPEATerrain

Cowpea terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Cowpea.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

COWPEAWater

Cowpea water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Cowpea.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv4 - Relative humidity at harvest stage (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CUCUMBERSoil

Cucumber soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Cucumber.

Format

A data frame with 10 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CUCUMBERTemp

Cucumber temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Cucumber.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CUCUMBERTerrain *Cucumber terrain requirement for land evaluation*

Description

A dataset containing the terrain characteristics of the crop requirements for farming Cucumber.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- SlopeD - Slope (degree, 6 classes)
- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

CUCUMBERWater *Cucumber water requirement for land evaluation*

Description

A dataset containing the water characteristics of the crop requirements for farming Cucumber.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

GROUNDNUTSSoil

Groundnuts soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Groundnuts.

Format

A data frame with 15 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECs - Apparent CEC Soil (cmol (+)/kg soil)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- OC - Organic carbon (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

GROUNDNUTSTemp

Groundnuts temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Groundnuts.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMaxAv - Mean max temp. of growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

GROUNDNUTSTerrain

Groundnuts terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Groundnuts.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 GROUNDNUTSWater

Groundnuts water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Groundnuts.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 GUAVASoil

Guava soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Guava.

Format

A data frame with 9 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

GUAVATemp

Guava temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Guava.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 GUAVATerrain

Guava terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Guava.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 GUAVAWater

Guava water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Guava.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

LaoCaiLT

Land and Terrain Characteristics of Lao Cai, Vietnam

Description

A dataset containing the land and terrain characteristics of the land units in Lao Cai, Vietnam.

Format

A data frame with 2928 rows and 10 columns

Details

- SlopeD - Slope (degree, 6 classes);
- CFragm - Coarse fragment (Vol.%);
- SoilDpt - Soil depth (cm);
- SoilTe - 12 classes of soil texture (Soil Taxonomy);
- CECc - Apparent CEC Clay (cmol (+)/kg clay);
- SumBCs - Sum of basic cations (cmol (+)/kg soil);
- pHH2O - pH H2O;
- BS - Base Saturation (%);
- OC - Organic carbon (%);
- Flood - Flooding;

LaoCaiTemp

Temperature Characteristics of Lao Cai, Vietnam

Description

A dataset containing the temperature characteristics of the land units in Lao Cai, Vietnam.

Format

A data frame with 2928 rows and 12 columns

Details

- Jan - January;
- Feb - February;
- Mar - March;
- Apr - April;
- May - May;
- Jun - June;
- Jul - July;
- Aug - August;
- Sep - September;
- Oct - October;
- Nov - November;
- Dec - December;

LaoCaiWater

Water Characteristics of Lao Cai, Vietnam

Description

A dataset containing the water characteristics of the land units in Lao Cai, Vietnam.

Format

A data frame with 2928 rows and 12 columns

Details

- Jan - January;
- Feb - February;
- Mar - March;
- Apr - April;
- May - May;
- Jun - June;
- Jul - July;
- Aug - August;
- Sep - September;
- Oct - October;
- Nov - November;
- Dec - December;

LONGANSoil

Longan soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Longan.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 LONGANTemp

Longan temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Longan.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TdMinXm - Mean daily minimum temperature of coldest month (°C)
- TyMinAv - Mean annual minimum temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 LONGANTerrain

Longan terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Longan.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

LONGANWater

Longan water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Longan.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MAIZESoil

Maize soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Maize.

Format

A data frame with 15 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- OC - Organic carbon (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MAIZETemp

Maize temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Maize.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MAIZETerrain

Maize terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Maize.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MAIZEWATER

Maize water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Maize.

Format

A data frame with 9 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv3 - Relative humidity of maturation Stage (%)
- WmnN2 - n/N develop. Stage (2nd month)
- WmnN4 - n/N maturation stage (4th month)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MANGOSoil

Mango soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Mango.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCc - Sum of basic cations (cmol (+)/kg of clay)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MANGOTemp

*Mango temp requirement for land evaluation***Description**

A dataset containing the temp characteristics of the crop requirements for farming Mango.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TyMinAv - Mean annual minimum temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MANGOTerrain

*Mango terrain requirement for land evaluation***Description**

A dataset containing the terrain characteristics of the crop requirements for farming Mango.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1) Basin furrow irrigation
- Slope2 - Slope (%) (2) Mechanized, high management level
- Slope3 - Slope (%) (3) Manual, low management level
- Flood - Flooding
- Drainage - Drainage
- SlopeD - 6 classes of slope (Degree)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MANGOWater

Mango water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Mango.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WmAvDry - Monthly precipitation during dry season (mm)
- WyhAv - Mean annual rel. humidity (%)
- WmnN4 - n/N maturation stage (4th month)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MarinduqueLT

Land and Terrain Characteristics of Marinduque, Philippines

Description

A dataset containing the land and terrain characteristics of the land units in Marinduque, Philippines.

Format

A data frame with 881 rows and 6 columns

Details

- Lat - Latitude of Land Units;
- Lon - Longitude of Land Units;
- CECc - Apparent CEC Clay (cmol (+)/kg clay);
- pHH2O - pH H2O;
- CFragm - Coarse fragment (Vol.%);
- SoilTe - 12 classes of soil texture (Soil Taxonomy);

MarinduqueTemp

Temperature Characteristics of Marinduque, Philippines

Description

A dataset containing the temperature characteristics of the land units in Marinduque, Philippines.

Format

A data frame with 881 rows and 14 columns

Details

- Lat - Latitude of Land Units;
- Lon - Longitude of Land Units;
- Jan - January;
- Feb - February;
- Mar - March;
- Apr - April;
- May - May;
- Jun - June;
- Jul - July;
- Aug - August;
- Sep - September;
- Oct - October;
- Nov - November;
- Dec - December;

MarinduqueWater

Water Characteristics of Marinduque, Philippines

Description

A dataset containing the water characteristics of the land units in Marinduque, Philippines.

Format

A data frame with 881 rows and 14 columns

Details

- Lat - Latitude of Land Units;
- Lon - Longitude of Land Units;
- Jan - January;
- Feb - February;
- Mar - March;
- Apr - April;
- May - May;
- Jun - June;
- Jul - July;
- Aug - August;
- Sep - September;
- Oct - October;
- Nov - November;
- Dec - December;

MILLETSSoil

Millets soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Millets.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MILLETSTemp

Millets temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Millets.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMaxAv - Mean max temp. of growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MILLETSTerrain

Millets terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Millets.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.
-

MILLETWater

Millets water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Millets.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv4 - Relative humidity at harvest stage (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

OILPALMSoil

Oil Palm soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Oil Palm.

Format

A data frame with 10 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm2 - Coarse fragment in depth (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 OILPALMTemp

Oil Palm temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Oil Palm.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TyMaxAv - Mean annual maximum temperature (°C)
- TdMinXm - Mean daily minimum temperature of coldest month (°C)
- TyAv - Mean annual temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 OILPALMTerrain

Oil Palm terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Oil Palm.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope - Slope
- Flood - Flooding
- Drainage4 - Drainage (4) Medium & fine textured soils
- Drainage5 - Drainage (5) Coarse textured soils

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

OILPALMWater

Oil Palm water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Oil Palm.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WynN - Mean annual n/N

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

OLIVESSoil

Olives soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Olives.

Format

A data frame with 10 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)

- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pH_{H2O} - pH H₂O
- OC - Organic carbon (%)
- EC_{dS} - EC_e (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

OLIVESTemp

Olives temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Olives.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- Ty_{Av} - Mean annual temperature (°C)
- Tm_{MinXm} - Average minimum temperature of coldest month (C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

OLIVESTerrain

Olives terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Olives.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

OLIVESWater

Olives water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Olives.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmSpecial1 - Monthly rainfall during the sclerification of stone (mm) - August (N hem) February (S hem.)
- WmSpecial2 - Monthly rainfall during the sclerification of stone (mm) - September (N hem) March (S hem.)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

ONIONSoil

Onion soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Onion.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

ONIONTemp

Onion temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Onion.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

ONIONTerrain

Onion terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Onion.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 ONIONWater

Onion water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Onion.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- TmAvDlen3 - Daylength (h) during yield form. Period

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 overall_suit

Overall Suitability Scores/Class of the Land Units

Description

This function computes the overall suitability scores and class of the land units.

Usage

```
overall_suit(suit, method = NULL, interval = NULL)
```

Arguments

suit	an object of class suitability.
method	a character for the method for computing the overall suitability, choices are: "minimum", "maximum", and "average". If NULL, method is set to "minimum".
interval	if NULL, the interval of the suitability class are the following: 0% - 25% (Not suitable, N), 25% - 50% (Marginally Suitable, S3), 50% - 75% (Moderately Suitable, S2), and 75% - 100% (Highly Suitable, S1). But users can assign custom intervals by specifying the values of the end points of the intervals. Say for intervals: 0% - 20% (Not suitable, N), 20% - 50% (Marginally Suitable, S3), 50% - 80% (Moderately Suitable, S2), and 80% - 100% (Highly Suitable, S1), is equivalent to interval = c(0, 0.2, 0.5, 0.8, 1).

Value

A data frame with columns:

- Score - the overall suitability scores
- Class - the overall suitability classes

See Also

[suit](https://alstat.github.io/ALUES/), <https://alstat.github.io/ALUES/>

Examples

```
library(ALUES)
out <- suit("ricebr", terrain=MarinduqueLT, water=MarinduqueWater, temp=MarinduqueTemp, sow_month=1)
out[["terrain"]]

# Soil Overall Suitability
head(overall_suit(out[["soil"]]))
head(overall_suit(out[["soil"]], "average"))
head(overall_suit(out[["soil"]], "maximum"))
head(overall_suit(out[["soil"]], "average", c(0, 0.3, 0.35, 0.6, 1.0)))

# Water Overall Suitability
head(overall_suit(out[["water"]], "average"))
head(overall_suit(out[["water"]], "maximum"))
head(overall_suit(out[["water"]], "average", c(0, 0.3, 0.35, 0.6, 1.0)))

# Temperature Overall Suitability
head(overall_suit(out[["temp"]], "average"))
head(overall_suit(out[["temp"]], "maximum"))
head(overall_suit(out[["temp"]], "average", c(0, 0.3, 0.35, 0.6, 1.0)))
```

PAPAYASoil

Papaya soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Papaya.

Format

A data frame with 9 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragn - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PAPAYATemp*Papaya temp requirement for land evaluation*

Description

A dataset containing the temp characteristics of the crop requirements for farming Papaya.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TyMinAv - Mean annual minimum temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PAPAYATerrain

Papaya terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Papaya.

Format

A data frame with 0 rows and 8 columns

Details

The following are the factors for evaluation:

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PAPAYAWater

Papaya water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Papaya.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WynN - Mean annual n/N
- WyhAv - Mean annual rel. humidity (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEACHSoil

Peach soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Peach.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 PEACHTemp

Peach temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Peach.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 PEACHTerrain

Peach terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Peach.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEACHWater

Peach water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Peach.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEARSoil

Pear soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Pear.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)

- pH_{H2O} - pH H₂O
- OC - Organic carbon (%)
- EC_{dS} - EC_e (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEARTemp

Pear temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Pear.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEARTerrain

Pear terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Pear.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEARWater

Pear water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Pear.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEASoil

Pea soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Pea.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragn - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEATemp

Pea temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Pea.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMinAv0 - Mean min. temp. at germination (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEATerrain

Pea terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Pea.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEAWater

Pea water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Pea.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEPPERGRSoil

Green Pepper soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Green Pepper.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic caions (cmol (+)/kg soil)

- pH_{H2O} - pH H₂O
- OC - Organic carbon (%)
- EC_{dS} - EC_e (dS/m)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEPPERGRTemp

Green Pepper temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Green Pepper.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEPPERGRTerrain

Green Pepper terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Green Pepper.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- Slope - nan

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PEPPERGRWater

Green Pepper water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Green Pepper.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PERSIMMONSoil

Persimmon soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Persimmon.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PERSIMMONTemp	<i>Persimmon temp requirement for land evaluation</i>
---------------	---

Description

A dataset containing the temp characteristics of the crop requirements for farming Persimmon.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PERSIMMONTerrain	<i>Persimmon terrain requirement for land evaluation</i>
------------------	--

Description

A dataset containing the terrain characteristics of the crop requirements for farming Persimmon.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PERSIMMONWater *Persimmon water requirement for land evaluation*

Description

A dataset containing the water characteristics of the crop requirements for farming Persimmon.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PINEAPPLESoil *Pineapple soil requirement for land evaluation*

Description

A dataset containing the soil characteristics of the crop requirements for farming Pineapple.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic caions (cmol (+)/kg soil)

- pH_{H2O} - pH H₂O
- OC - Organic carbon (%)
- EC_{dS} - EC_e (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PINEAPPLETemp

Pineapple temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Pineapple.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PINEAPPLETerrain

Pineapple terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Pineapple.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PINEAPPLEWater

Pineapple water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Pineapple.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WyhAv - Mean annual rel. humidity (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PLUMSoil

Plum soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Plum.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PLUMTemp *Plum temp requirement for land evaluation*

Description

A dataset containing the temp characteristics of the crop requirements for farming Plum.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PLUMTerrain *Plum terrain requirement for land evaluation*

Description

A dataset containing the terrain characteristics of the crop requirements for farming Plum.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

PLUMWater

Plum water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Plum.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

POTATOSoil

Potato soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Potato.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt1 - Surface Soil Depth (cm)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)

- pH_{H2O} - pH H₂O
- OC - Organic carbon (%)
- EC_{dS} - EC_e (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

POTATOSWSoil

Sweet Potato soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Sweet Potato.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO₃ - CaCO₃ (%)
- Gyps - Gypsum (%)
- CEC_c - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pH_{H2O} - pH H₂O
- OC - Organic carbon (%)
- EC_{dS} - EC_e (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 POTATOSWTemp

Sweet Potato temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Sweet Potato.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 POTATOSWTerrain

Sweet Potato terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Sweet Potato.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 POTATOSWater

Sweet Potato water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Sweet Potato.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WmhAv4 - Relative humidity at harvest stage (%)
- WmnN2 - n/N develop. Stage (2nd month)
- WmnN4 - n/N maturation stage (4th month)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 POTATOTemp

Potato temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Potato.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMinAv1 - Average absolut Min. temperature of the first month (°C)
- TmMinAv4 - Average absolut Min. temperature of other months (°C)
- TdAvgDiff - Average Temperature difference between day-night (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

POTATOTerrain

Potato terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Potato.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

POTATOWater

Potato water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Potato.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- TgAvDlen - Average daylength growing cycle (h)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICEBRSoil

Rainfed Bunded Rice soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Rainfed Bunded Rice.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICEBRTemp

Rainfed Bunded Rice temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Rainfed Bunded Rice.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmMinAv4 - Average absolut Min. temperature of other months (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICEBRTerrain

Rainfed Bunded Rice terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Rainfed Bunded Rice.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Flood - Flooding
- Drainage - Drainage

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICEBRWater

Rainfed Bunded Rice water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Rainfed Bunded Rice.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)

- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv4 - Relative humidity at harvest stage (%)
- WynN - Mean annual n/N

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICEIWSoil

Irrigated Rice soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Irrigated Rice.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- pHH2O - pH H2O
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 RICEIWTemp

Irrigated Rice temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Irrigated Rice.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmMinAv4 - Average absolut Min. temperature of other months (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 RICEIWTerrain

Irrigated Rice terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Irrigated Rice.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICEIWWater

Irrigated Rice water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Irrigated Rice.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv4 - Relative humidity at harvest stage (%)
- WynN - Mean annual n/N

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICENFSoil	<i>Rice Cultivation under Natural Floods soil requirement for land evaluation</i>
------------	---

Description

A dataset containing the soil characteristics of the crop requirements for farming Rice Cultivation under Natural Floods.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICENFTemp	<i>Rice Cultivation under Natural Floods temp requirement for land evaluation</i>
------------	---

Description

A dataset containing the temp characteristics of the crop requirements for farming Rice Cultivation under Natural Floods.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmMinAv4 - Average absolut Min. temperature of other months (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICENFTerrain	<i>Rice Cultivation under Natural Floods terrain requirement for land evaluation</i>
---------------	--

Description

A dataset containing the terrain characteristics of the crop requirements for farming Rice Cultivation under Natural Floods.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- Flood - Flooding
- Drainage - Drainage

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICENFWater

Rice Cultivation under Natural Floods water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Rice Cultivation under Natural Floods.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv4 - Relative humidity at harvest stage (%)
- WynN - Mean annual n/N

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICEURSoil

Rainfed Upland Rice soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Rainfed Upland Rice.

Format

A data frame with 8 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICEURTemp

Rainfed Upland Rice temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Rainfed Upland Rice.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmMinAv4 - Average absolut Min. temperature of other months (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICEURTerrain

Rainfed Upland Rice terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Rainfed Upland Rice.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RICEURWater

Rainfed Upland Rice water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Rainfed Upland Rice.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv4 - Relative humidity at harvest stage (%)
- WynN - Mean annual n/N

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RUBBERSoil

Rubber soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Rubber.

Format

A data frame with 10 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RUBBERTemp

Rubber temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Rubber.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TyMaxAv - Mean annual maximum temperature (°C)
- TdMinXm - Mean daily minimum temperature of coldest month (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RUBBERTerrain	<i>Rubber terrain requirement for land evaluation</i>
---------------	---

Description

A dataset containing the terrain characteristics of the crop requirements for farming Rubber.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Flood - Flooding
- Drainage - Drainage
- Slope - nan

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

RUBBERWater	<i>Rubber water requirement for land evaluation</i>
-------------	---

Description

A dataset containing the water characteristics of the crop requirements for farming Rubber.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmER - Months of excessive rain (x)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SAFFLOWERSoil

Safflower soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Safflower.

Format

A data frame with 8 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.
-

SAFFLOWERTemp

Safflower temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Safflower.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TmAv1 - Mean temp. of the initial stage(C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SAFFLOWERTerrain

Safflower terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Safflower.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SAFFLOWERWater

Safflower water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Safflower.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv3 - Relative humidity of maturation Stage (%)
- WmhAv4 - Relative humidity at harvest stage (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SESAMESoil

Sesame soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Sesame.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SESAMETemp

Sesame temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Sesame.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMaxAv - Mean max temp. of growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SESAMETerrain

Sesame terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Sesame.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 SESAMEWater

Sesame water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Sesame.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 SORGHUMSoil

Sorghum soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Sorghum.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SORGHUMTemp

Sorghum temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Sorghum.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMaxAv - Mean max temp. of growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SORGHUMTerrain *Sorghum terrain requirement for land evaluation*

Description

A dataset containing the terrain characteristics of the crop requirements for farming Sorghum.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SORGHUMWater *Sorghum water requirement for land evaluation*

Description

A dataset containing the water characteristics of the crop requirements for farming Sorghum.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SOYASoil

Soya soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Soya.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 SOYATemp

Soya temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Soya.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 SOYATerrain

Soya terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Soya.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SOYAWater

Soya water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Soya.

Format

A data frame with 9 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv3 - Relative humidity of maturation Stage (%)
- WmnN2 - n/N develop. Stage (2nd month)
- WmnN4 - n/N maturation stage (4th month)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SUGARCANESoil

Sugar Cane soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Sugar Cane.

Format

A data frame with 15 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- OC - Organic carbon (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 SUGARCANETemp

Sugar Cane temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Sugar Cane.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- TdAvg0 - Mean day temperature at germination stage(°C)
- TdAvg1 - Mean day temperature for tillage stage (°C)
- TdAvg2 - Mean day temperature for vegetative stage (°C)
- Tcoef - $(T_{max}-T_{min})/T_{mean}$ maturation stage

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 SUGARCANETerrain

Sugar Cane terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Sugar Cane.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 SUGARCANEWater

Sugar Cane water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Sugar Cane.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- Wd10 - 10 days of rainfall (mm)
- SunH - Sunshine : hours/year
- WynN - Mean annual n/N
- WmhAv3 - Relative humidity of maturation Stage (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 suit

Suitability Scores/Class of the Land Units

Description

This function calculates the suitability scores and class of the land units.

Usage

```

suit(
  crop,
  terrain = NULL,
  water = NULL,
  temp = NULL,
  mf = "triangular",
  sow_month = NULL,
  minimum = NULL,
  maximum = "average",
  interval = NULL,
  sigma = NULL
)

```

Arguments

crop	a string for the name of the crop;
terrain	a data frame for the terrain characteristics of the input land units;
water	a data frame for the water characteristics of the input land units;
temp	a data frame for the temperature characteristics of the input land units;
mf	membership function with default assigned to "triangular" fuzzy model. Other fuzzy models included are "trapezoidal" and "gaussian".
sow_month	sowing month of the crop. Takes integers from 1 to 12 (inclusive), representing the twelve months of the year. So if sets to 1, the function assumes sowing month to be January.
minimum	factor's minimum value. If NULL (default), minimum is set to 0. But if numeric of length one, say 0.5, then minimum is set to 0.5, for all factors. To set multiple minimums for multiple factors, simply concatenate these into a numeric vector, the length of this vector should be equal to the number of factors in input land units parameters. However, it can also be set to "average", please refer to the online documentation for more, link in the "See Also" section below.
maximum	maximum value for factors. To set multiple maximums for multiple factors, simply concatenate these into a numeric vector, the length of this vector should be equal to the number of factors in input land units parameters. However, it can also be set to "average", please refer to the online documentation for more, link in the "See Also" section below.
interval	domains for every suitability class (S1, S2, S3). If fixed (NULL), the interval would be 0 to 25% for N (Not Suitable), 25% to 50% for S3 (Marginally Suitable), 50% to 75% for S2 (Moderately Suitable), and 75% to 100% for (Highly Suitable). If "unbias", the package will take into account the shape of the membership function, and provide the appropriate suitability class intervals. However, it can also be customized by specifying the limits of the suitability classes. Please refer to the online documentation for more, link in the "See Also" section below.
sigma	If mf = "gaussian", then sigma represents the constant sigma in the Gaussian formula.

Value

A list of outputs of target characteristics, with the following components:

- "terrain" - a list of outputs for terrain characteristics
- "soil" - a list of outputs for soil characteristics
- "water" - a list of outputs for water characteristics
- "temp" - a list of outputs for temperature characteristics

These components are only available when specified as the target characteristics in either of the arguments above, that is, if terrain argument is specified above, then the "terrain" and "soil" components will be available in the output list. This is also true if water and temp are specified in the arguments above.

Each of the components returned above contains a list of outputs as well with the following components:

- "Factors Evaluated" - a character of factors that matched between the input land units factor and the targetted crop requirement factor
- "Suitability Score" - a data frame of suitability scores for each of the matched factors
- "Suitability Class" - a data frame of suitability classes for each of the matched factors
- "Factors' Minimum Values" - a numeric of minimum values used in the membership function for computing the suitability scores
- "Factors' Maximum Values" - a numeric of maximum values used in the membership function for computing the suitability scores
- "Factors' Weights" - a numeric of weights of the factors specified in the input crop requirements
- "Crop Evaluated" - a character of the name of the targetted crop requirement dataset

See Also

<https://alstat.github.io/ALUES/>

Examples

```
library(ALUES)

rice_suit <- suit("ricebr", water=MarinduqueWater, temp=MarinduqueTemp, sow_month = 1)
lapply(rice_suit[["water"]], function(x) head(x)) # access results for water suitability
lapply(rice_suit[["temp"]], function(x) head(x)) # access results for temperature suitability
rice_suit <- suit("ricebr", terrain=MarinduqueLT)
lapply(rice_suit[["terrain"]], function(x) head(x))
lapply(rice_suit[["soil"]], function(x) head(x))
```

 suitability

Suitability Scores/Class of the Land Units

Description

This function calculates the suitability scores and class of the land units.

Usage

```
suitability(
  x,
  y,
  mf = "triangular",
  sow_month = NULL,
  minimum = NULL,
  maximum = "average",
  interval = NULL,
  sigma = NULL
)
```

Arguments

x	a data frame consisting the properties of the land units;
y	a data frame consisting the requirements of a given characteristics (terrain, soil, water and temperature) for a given crop (e.g. coconut, cassava, etc.);
mf	membership function with default assigned to "triangular" fuzzy model. Other fuzzy models included are "trapezoidal" and "gaussian".
sow_month	sowing month of the crop. Takes integers from 1 to 12 (inclusive), representing the twelve months of the year. So if sets to 1, the function assumes sowing month to be January.
minimum	factor's minimum value. If NULL (default), minimum is set to 0. But if numeric of length one, say 0.5, then minimum is set to 0.5, for all factors. To set multiple minimums for multiple factors, simply concatenate these into a numeric vector, the length of this vector should be equal to the number of factors in input land units parameters. However, it can also be set to "average", please refer to the online documentation for more, link in the "See Also" section below.
maximum	maximum value for factors. To set multiple maximums for multiple factors, simply concatenate these into a numeric vector, the length of this vector should be equal to the number of factors in input land units parameters. However, it can also be set to "average", please refer to the online documentation for more, link in the "See Also" section below.
interval	domains for every suitability class (S1, S2, S3). If fixed (NULL), the interval would be 0 to 25% for N (Not Suitable), 25% to 50% for S3 (Marginally Suitable), 50% to 75% for S2 (Moderately Suitable), and 75% to 100% for (Highly

Suitable). If "unbias", the package will take into account the shape of the membership function, and provide the appropriate suitability class intervals. However, it can also be customized by specifying the limits of the suitability classes. Please refer to the online documentation for more, link in the "See Also" section below.

sigma If mf = "gaussian", then sigma represents the constant sigma in the Gaussian formula.

Value

A list with the following components:

- "Factors Evaluated" - a character of factors that matched between the input land units factor and the targetted crop requirement factor
- "Suitability Score" - a data frame of suitability scores for each of the matched factors
- "Suitability Class" - a data frame of suitability classes for each of the matched factors
- "Factors' Minimum Values" - a numeric of minimum values used in the membership function for computing the suitability scores
- "Factors' Maximum Values" - a numeric of maximum values used in the membership function for computing the suitability scores
- "Factors' Weights" - a numeric of weights of the factors specified in the input crop requirements
- "Crop Evaluated" - a character of the name of the targetted crop requirement dataset

#' @seealso <https://alstat.github.io/ALUES/>

SUNFLOWERSoil

Sunflower soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Sunflower.

Format

A data frame with 11 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)

- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pH_{H2O} - pH H₂O
- OC - Organic carbon (%)
- EC_{dS} - EC_e (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SUNFLOWERTemp

Sunflower temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Sunflower.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- Tg_{Av} - Mean temperature of the growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SUNFLOWERTerrain *Sunflower terrain requirement for land evaluation*

Description

A dataset containing the terrain characteristics of the crop requirements for farming Sunflower.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

SUNFLOWERWater *Sunflower water requirement for land evaluation*

Description

A dataset containing the water characteristics of the crop requirements for farming Sunflower.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmAv5 - Mean precipitation of fifth month (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 TEASoil

Tea soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Tea.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 TEATemp

Tea temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Tea.

Format

A data frame with 5 rows and 8 columns

Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TmMinAv - Mean min. temp. of warmest month (°C)
- TmMinXm - Avarage minimum temperature of coldest month (C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv4Xm - Mean temp. of 4 warmest month (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 TEATerrain

Tea terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Tea.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

TEAWater

Tea water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Tea.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WmDryLen - Length dry season (months : $P < 1/2$ PET)
- WyhAv - Mean annual rel. humidity (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

TOBACCOSoil

Tobacco soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Tobacco.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 TOBACCOTemp

Tobacco temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Tobacco.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 TOBACCOTerrain

Tobacco terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Tobacco.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 TOBACCOWater

Tobacco water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Tobacco.

Format

A data frame with 3 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WynN - Mean annual n/N
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

 TOMATOSoil

Tomato soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Tomato.

Format

A data frame with 12 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

TOMATOTemp

Tomato temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Tomato.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)
- TmAv3 - Mean temp. of the flowering stage (°C)
- TdDiff3 - Temp. diff day/night flowering stage (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

TOMATOTerrain *Tomato terrain requirement for land evaluation*

Description

A dataset containing the terrain characteristics of the crop requirements for farming Tomato.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

TOMATOWater *Tomato water requirement for land evaluation*

Description

A dataset containing the water characteristics of the crop requirements for farming Tomato.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

WATERMELONSoil

Watermelon soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Watermelon.

Format

A data frame with 10 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - nan

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

WATERMELONTemp

Watermelon temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Watermelon.

Format

A data frame with 1 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.
-

WATERMELONTerrain

Watermelon terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Watermelon.

Format

A data frame with 7 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization.)
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

WATERMELONWater

Watermelon water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Watermelon.

Format

A data frame with 2 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

WHEATSoil

Wheat soil requirement for land evaluation

Description

A dataset containing the soil characteristics of the crop requirements for farming Wheat.

Format

A data frame with 14 rows and 8 columns

Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

WHEATTemp

Wheat temp requirement for land evaluation

Description

A dataset containing the temp characteristics of the crop requirements for farming Wheat.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmAv3 - Mean temp. of the flowering stage (°C)
- TmAv4 - Mean temp. of the ripening stage (°C)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

WHEATTerrain

Wheat terrain requirement for land evaluation

Description

A dataset containing the terrain characteristics of the crop requirements for farming Wheat.

Format

A data frame with 6 rows and 8 columns

Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization.)
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

WHEATWater

Wheat water requirement for land evaluation

Description

A dataset containing the water characteristics of the crop requirements for farming Wheat.

Format

A data frame with 4 rows and 8 columns

Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)

See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

Index

* dataset

ALFALFASoil, 7
ALFALFATemp, 8
ALFALFATerrain, 9
ALFALFAWater, 9
AVOCADOSoil, 10
AVOCADOTemp, 11
AVOCADOTerrain, 11
AVOCADOWater, 12
BAMBOOSoil, 12
BAMBOOTemp, 13
BAMBOOTerrain, 13
BAMBOOWater, 14
BANANASoil, 14
BANANATemp, 15
BANANATerrain, 16
BANANAWater, 16
BARLEYSoil, 17
BARLEYTemp, 18
BARLEYTerrain, 18
BARLEYWater, 19
BEANCAsoil, 19
BEANCATemp, 20
BEANCATerrain, 21
BEANCAWater, 21
BEANSSoil, 22
BEANSTemp, 23
BEANSTerrain, 23
BEANSWater, 24
CABBAGESoil, 24
CABBAGETemp, 25
CABBAGETerrain, 26
CABBAGEWater, 26
CARROTSSoil, 27
CARROTSTemp, 28
CARROTSTerrain, 28
CARROTSWater, 29
CASHEWSoil, 29
CASHEWTemp, 30
CASHEWTerrain, 31
CASHEWWater, 31
CASSAVASoil, 32
CASSAVATemp, 33
CASSAVATerrain, 33
CASSAVAWater, 34
CHICKPEASoil, 34
CHICKPEATemp, 35
CHICKPEATerrain, 36
CHICKPEAWater, 36
CINNAMONSoil, 37
CINNAMONTemp, 38
CINNAMONTerrain, 38
CINNAMONWater, 39
CITRUSSoil, 39
CITRUSTemp, 40
CITRUSTerrain, 41
CITRUSWater, 41
COCOASoil, 42
COCOATemp, 43
COCOATerrain, 43
COCOAWater, 44
COCONUTSoil, 44
COCONUTTemp, 45
COCONUTTerrain, 46
COCONUTWater, 46
COFFEEARSoil, 47
COFFEEARTemp, 48
COFFEEARTerrain, 48
COFFEEARWater, 49
COFFEEEROSoil, 49
COFFEEEROTemp, 50
COFFEEEROTerrain, 51
COFFEEEROWater, 51
COTTONSoil, 52
COTTONTemp, 53
COTTONTerrain, 53
COTTONWater, 54
COWPEASoil, 55

- COWPEATemp, [55](#)
- COWPEATerrain, [56](#)
- COWPEAWater, [57](#)
- CUCUMBERSoil, [57](#)
- CUCUMBERTemp, [58](#)
- CUCUMBERTerrain, [59](#)
- CUCUMBERWater, [59](#)
- GROUNDNUTSoil, [60](#)
- GROUNDNUTSTemp, [61](#)
- GROUNDNUTSTerrain, [61](#)
- GROUNDNUTSWater, [62](#)
- GUAVASoil, [62](#)
- GUAVATemp, [63](#)
- GUAVATerrain, [64](#)
- GUAVAWater, [64](#)
- LaoCaiLT, [65](#)
- LaoCaiTemp, [65](#)
- LaoCaiWater, [66](#)
- LONGANSoil, [67](#)
- LONGANTemp, [68](#)
- LONGANTerrain, [68](#)
- LONGANWater, [69](#)
- MAIZESoil, [69](#)
- MAIZETemp, [70](#)
- MAIZETerrain, [71](#)
- MAIZEWater, [71](#)
- MANGOSoil, [72](#)
- MANGOTemp, [73](#)
- MANGOTerrain, [73](#)
- MANGOWater, [74](#)
- MarinduqueLT, [74](#)
- MarinduqueTemp, [75](#)
- MarinduqueWater, [76](#)
- MILLETSoil, [76](#)
- MILLETSTemp, [77](#)
- MILLETSTerrain, [78](#)
- MILLETSWater, [78](#)
- OILPALMSoil, [79](#)
- OILPALMTemp, [80](#)
- OILPALMTerrain, [80](#)
- OILPALMWater, [81](#)
- OLIVESSoil, [81](#)
- OLIVESTemp, [82](#)
- OLIVESTerrain, [83](#)
- OLIVESWater, [83](#)
- ONIONSoil, [84](#)
- ONIONTemp, [85](#)
- ONIONTerrain, [85](#)
- ONIONWater, [86](#)
- PAPAYASoil, [88](#)
- PAPAYATemp, [88](#)
- PAPAYATerrain, [89](#)
- PAPAYAWater, [89](#)
- PEACHSoil, [90](#)
- PEACHTemp, [91](#)
- PEACHTerrain, [91](#)
- PEACHWater, [92](#)
- PEARSoil, [92](#)
- PEARTemp, [93](#)
- PEARTerrain, [93](#)
- PEARWater, [94](#)
- PEASoil, [95](#)
- PEATemp, [95](#)
- PEATerrain, [96](#)
- PEAWater, [97](#)
- PEPPERGRSoil, [97](#)
- PEPPERGRTemp, [98](#)
- PEPPERGRTerrain, [98](#)
- PEPPERGRWater, [99](#)
- PERSIMMONSoil, [100](#)
- PERSIMMONTemp, [101](#)
- PERSIMMONTerrain, [101](#)
- PERSIMMONWater, [102](#)
- PINEAPPLESoil, [102](#)
- PINEAPPLETemp, [103](#)
- PINEAPPLETerrain, [103](#)
- PINEAPPLEWater, [104](#)
- PLUMSoil, [105](#)
- PLUMTemp, [106](#)
- PLUMTerrain, [106](#)
- PLUMWater, [107](#)
- POTATOSoil, [107](#)
- POTATOSWSoil, [108](#)
- POTATOSWTemp, [109](#)
- POTATOSWTerrain, [109](#)
- POTATOSWWater, [110](#)
- POTATOTemp, [110](#)
- POTATOTerrain, [111](#)
- POTATOWater, [112](#)
- RICEBRSoil, [112](#)
- RICEBRTemp, [113](#)
- RICEBRTerrain, [114](#)
- RICEBRWater, [114](#)
- RICEIWSoil, [115](#)
- RICEIWTemp, [116](#)
- RICEIWTerrain, [116](#)

- RICEIWater, 117
- RICENSoil, 118
- RICENTemp, 119
- RICENTerrain, 119
- RICENWater, 120
- RICEURSoil, 121
- RICEURTemp, 121
- RICEURTerrain, 122
- RICEURWater, 123
- RUBBERSoil, 123
- RUBBERTemp, 124
- RUBBERTerrain, 125
- RUBBERWater, 125
- SAFFLOWERSoil, 126
- SAFFLOWERTemp, 126
- SAFFLOWERTerrain, 127
- SAFFLOWERWater, 127
- SESAMESoil, 128
- SESAMETemp, 129
- SESAMETerrain, 129
- SESAMEWater, 130
- SORGHUMSoil, 130
- SORGHUMTemp, 131
- SORGHUMTerrain, 132
- SORGHUMWater, 132
- SOYASoil, 133
- SOYATemp, 134
- SOYATerrain, 134
- SOYAWater, 135
- SUGARCANESoil, 136
- SUGARCANETemp, 137
- SUGARCANETerrain, 137
- SUGARCANEWater, 138
- SUNFLOWERSoil, 142
- SUNFLOWERTemp, 143
- SUNFLOWERTerrain, 144
- SUNFLOWERWater, 144
- TEASoil, 145
- TEATemp, 146
- TEATerrain, 146
- TEAWater, 147
- TOBACCSoil, 148
- TOBACCOTemp, 149
- TOBACCOTerrain, 149
- TOBACOWater, 150
- TOMATOSoil, 150
- TOMATOTemp, 151
- TOMATOTerrain, 152
- TOMATOWater, 152
- WATERMELONSoil, 153
- WATERMELONTemp, 154
- WATERMELONTerrain, 154
- WATERMELONWater, 155
- WHEATSoil, 155
- WHEATTemp, 156
- WHEATTerrain, 157
- WHEATWater, 157

- ALFALFASoil, 7
- ALFALFATemp, 8
- ALFALFATerrain, 9
- ALFALFAWater, 9
- ALUES-package, 7
- AVOCADOSoil, 10
- AVOCADOTemp, 11
- AVOCADOTerrain, 11
- AVOCADOWater, 12

- BAMBOOSoil, 12
- BAMBOOTemp, 13
- BAMBOOTerrain, 13
- BAMBOOWater, 14
- BANANASoil, 14
- BANANATemp, 15
- BANANATerrain, 16
- BANANAWater, 16
- BARLEYSoil, 17
- BARLEYTemp, 18
- BARLEYTerrain, 18
- BARLEYWater, 19
- BEANCASoil, 19
- BEANCATemp, 20
- BEANCATerrain, 21
- BEANCAWater, 21
- BEANSSoil, 22
- BEANSTemp, 23
- BEANSTerrain, 23
- BEANSWater, 24

- CABBAGESoil, 24
- CABBAGETemp, 25
- CABBAGETerrain, 26
- CABBAGEWater, 26
- CARROTSSoil, 27
- CARROTSTemp, 28
- CARROTSTerrain, 28
- CARROTSWater, 29

- CASHEWSoil, 29
 CASHEWTemp, 30
 CASHEWTerrain, 31
 CASHEWWater, 31
 CASSAVASoil, 32
 CASSAVATemp, 33
 CASSAVATerrain, 33
 CASSAVAWater, 34
 CHICKPEASoil, 34
 CHICKPEATemp, 35
 CHICKPEATerrain, 36
 CHICKPEAWater, 36
 CINNAMONSoil, 37
 CINNAMONTemp, 38
 CINNAMONTerrain, 38
 CINNAMONWater, 39
 CITRUSSoil, 39
 CITRUSTemp, 40
 CITRUSTerrain, 41
 CITRUSWater, 41
 COCOASoil, 42
 COCOATemp, 43
 COCOATerrain, 43
 COCOAWater, 44
 COCONUTSoil, 44
 COCONUTTemp, 45
 COCONUTTerrain, 46
 COCONUTWater, 46
 COFFEEARSoil, 47
 COFFEEARTemp, 48
 COFFEEARTerrain, 48
 COFFEEARWater, 49
 COFFEEROSoil, 49
 COFFEEROTemp, 50
 COFFEEROTerrain, 51
 COFFEEROWater, 51
 COTTONSoil, 52
 COTTONTemp, 53
 COTTONTerrain, 53
 COTTONWater, 54
 COWPEASoil, 55
 COWPEATemp, 55
 COWPEATerrain, 56
 COWPEAWater, 57
 CUCUMBERSoil, 57
 CUCUMBERTemp, 58
 CUCUMBERTerrain, 59
 CUCUMBERWater, 59
 GROUNDNUTSoil, 60
 GROUNDNUTTemp, 61
 GROUNDNUTTerrain, 61
 GROUNDNUTWater, 62
 GUAVASoil, 62
 GUAVATemp, 63
 GUAVATerrain, 64
 GUAVAWater, 64
 LaoCaiLT, 65
 LaoCaiTemp, 65
 LaoCaiWater, 66
 LONGANSoil, 67
 LONGANTemp, 68
 LONGANTerrain, 68
 LONGANWater, 69
 MAIZESoil, 69
 MAIZETemp, 70
 MAIZETerrain, 71
 MAIZEWater, 71
 MANGOSoil, 72
 MANGOTemp, 73
 MANGOTerrain, 73
 MANGOWater, 74
 MarinduqueLT, 74
 MarinduqueTemp, 75
 MarinduqueWater, 76
 MILLETSoil, 76
 MILLETSTemp, 77
 MILLETSTerrain, 78
 MILLETWater, 78
 OILPALMSoil, 79
 OILPALMTemp, 80
 OILPALMTerrain, 80
 OILPALMWater, 81
 OLIVESSoil, 81
 OLIVESTemp, 82
 OLIVESTerrain, 83
 OLIVESWater, 83
 ONIONSoil, 84
 ONIONTemp, 85
 ONIONTerrain, 85
 ONIONWater, 86
 overall_suit, 86
 PAPAYASoil, 88
 PAPAYATemp, 88

- PAPAYATerrain, 89
 PAPAYAWater, 89
 PEACHSoil, 90
 PEACHTemp, 91
 PEACHTerrain, 91
 PEACHWater, 92
 PEARSoil, 92
 PEARTemp, 93
 PEARTerrain, 93
 PEARWater, 94
 PEASoil, 95
 PEATemp, 95
 PEATerrain, 96
 PEAWater, 97
 PEPPERGRSoil, 97
 PEPPERGRTemp, 98
 PEPPERGRTerrain, 98
 PEPPERGRWater, 99
 PERSIMMONSoil, 100
 PERSIMMONTemp, 101
 PERSIMMONTerrain, 101
 PERSIMMONWater, 102
 PINEAPPLESoil, 102
 PINEAPPLETemp, 103
 PINEAPPLETerrain, 103
 PINEAPPLEWater, 104
 PLUMSoil, 105
 PLUMTemp, 106
 PLUMTerrain, 106
 PLUMWater, 107
 POTATOSoil, 107
 POTATOSWSoil, 108
 POTATOSWTemp, 109
 POTATOSWTerrain, 109
 POTATOSWWater, 110
 POTATOTemp, 110
 POTATOTerrain, 111
 POTATOWater, 112
- RICEBRSoil, 112
 RICEBRTemp, 113
 RICEBRTerrain, 114
 RICEBRWater, 114
 RICEIWSoil, 115
 RICEIWTemp, 116
 RICEIWTerrain, 116
 RICEIWWater, 117
 RICENFSoil, 118
 RICENFTemp, 119
- RICENFTerrain, 119
 RICENFWater, 120
 RICEURSoil, 121
 RICEURTemp, 121
 RICEURTerrain, 122
 RICEURWater, 123
 RUBBERSoil, 123
 RUBBERTemp, 124
 RUBBERTerrain, 125
 RUBBERWater, 125
- SAFFLOWERSoil, 126
 SAFFLOWERTemp, 126
 SAFFLOWERTerrain, 127
 SAFFLOWERWater, 127
 SESAMESoil, 128
 SESAMETemp, 129
 SESAMETerrain, 129
 SESAMEWater, 130
 SORGHUMSoil, 130
 SORGHUMTemp, 131
 SORGHUMTerrain, 132
 SORGHUMWater, 132
 SOYASoil, 133
 SOYATemp, 134
 SOYATerrain, 134
 SOYAWater, 135
 SUGARCANESoil, 136
 SUGARCANETemp, 137
 SUGARCANETerrain, 137
 SUGARCANEWater, 138
 suit, 87, 138
 suitability, 141
 SUNFLOWERSoil, 142
 SUNFLOWERTemp, 143
 SUNFLOWERTerrain, 144
 SUNFLOWERWater, 144
- TEASoil, 145
 TEATemp, 146
 TEATerrain, 146
 TEAWater, 147
 TOBACCSOIL, 148
 TOBACCOTemp, 149
 TOBACCOTerrain, 149
 TOBACCOWater, 150
 TOMATOSoil, 150
 TOMATOTemp, 151
 TOMATOTerrain, 152

TOMATOWater, [152](#)

WATERMELONSoil, [153](#)

WATERMELONTemp, [154](#)

WATERMELONTerrain, [154](#)

WATERMELONWater, [155](#)

WHEATSoil, [155](#)

WHEATTemp, [156](#)

WHEATTerrain, [157](#)

WHEATWater, [157](#)