

FAiR Vignette

Ben Goodrich

January 12, 2009

This vignette is intended to be a quick reference that defines terms and illustrates how to use the pop-up menus. Large parts will not make a great deal of sense outside the introduction to SEFA and lexical optimization in Goodrich (2008a) and Goodrich (2008b).

Contents

1	Notation	3
2	Discrepancy functions in FAiR	4
2.1	Estimating Gamma	4
3	Using <code>make_manifest()</code>	5
4	Esoteric definitions	7
4.1	The factor contribution matrix	7
4.2	Generalized variance (GV) and effective variance (EV)	8
5	Important theorems	8
5.1	Howe (1955) and Koopmans and Reiersøl (1950)	8
5.2	Reiersøl (1950)	9
6	SEFA or CFA via <code>Factanal()</code>	9
6.1	Number of Factors	10
6.2	Main menu	10
6.2.1	Fix specific coefficients to particular values (second and eighth checkboxes)	11
6.2.2	Constrain coefficients to be equal to each other (fourth checkbox)	13
6.2.3	Change bounds on coefficients (third checkbox)	13
6.2.4	Important Note	15
6.2.5	Nondefault mapping rule and changing the requisite number of zeros (11th and 12th checkboxes)	15
6.2.6	Change bounds on correlations among primary factors (first checkbox)	17
6.2.7	Impose functional inequality restrictions	18

7	Exploratory Factor Analysis via <code>Factnal()</code> and <code>Rotate()</code>	26
7.1	Factor Extraction	26
7.2	Factor Transformation	26
7.2.1	Functional inequality restrictions	26
7.2.2	The ultimate analytic criterion	28
8	References	31

1 Notation

Some notation is provided so that the reader can refer back to it. For simplicity, I usually do not distinguish the notation for a population parameter from the notation for an estimate of that population parameter here. In particular, note that all models currently in FAiR utilize an “embedded correlation” parameterization.

- n is the number of outcome variables, which are indexed by j
- Σ is the $n \times n$ covariance matrix among outcomes in the population
- S is the $n \times n$ covariance matrix among outcomes in the sample
- $s = \text{vecs}(S)$ stacks the elements below and including the diagonal of S into a vector of length $n^* = 0.5n(n + 1)$
- $\Gamma = \text{acov}(s)$ is the asymptotic covariance of s
- r is the number of factors, which are indexed by p
- β is the $n \times r$ primary pattern matrix with one column for each factor
- Φ is the $r \times r$ correlation matrix among primary factors
- Λ is used to indicate the primary pattern matrix only if $\Phi = I_r$
- $R^* = \beta\Phi\beta'$ is the reduced correlation matrix with communalities along the diagonal, denoted by $h_j^2 = R_{jj}^*$
- Θ^2 is the $n \times n$ diagonal (for now) covariance matrix among unique factors such that $\Theta_{jj}^2 = 1 - h_j^2$
- $R = R^* + \Theta^2 = \beta\Phi\beta' + \Theta^2$ is the $n \times n$ reproduced correlation matrix with ones along the diagonal
- Ω is the $n \times n$ diagonal matrix of standard deviations of the manifest variables to be estimated generally
- $C = \Omega R \Omega = \Omega(\beta\Phi\beta' + \Theta^2)\Omega$ is the $n \times n$ reproduced covariance matrix and also the expectation of S
- $c = \text{vecs}(C)$ stacks the elements below and including the diagonal of C into a vector of length n^*
- $\Pi = \beta\Phi$ is the $n \times r$ primary structure matrix of correlations between outcome variables and factors
- $\mathcal{D} = \beta \times \Pi$ is the $n \times r$ factor contribution matrix, where \times indicates element-by-element multiplication
- $D = [\text{Diag}(\Phi^{-1})]^{-\frac{1}{2}}$ is the $r \times r$ correlation matrix between primary and reference factors
- $\Psi = D\Phi^{-1}D$ is the $r \times r$ correlation matrix among reference factors
- $\Upsilon = \beta D$ is the $n \times r$ reference structure matrix
- $\Phi = \check{\beta}\check{\Phi}\check{\beta}' + \check{\Theta}^2$ is the $\check{r} \times \check{r}$ primary factor correlation matrix as a function of second-order parameters

In general, the “ notation indicates that the parameter pertains to level 2 of the model (if a two-level model is specified). The R code in FAiR uses the same notational conventions at level 1, but uses different symbols at level 2. The primary pattern coefficient at level two is called `Delta` and the correlation matrix among factors at level two is called `Xi`.

2 Discrepancy functions in FAiR

This section defines the discrepancy functions that can be specified in the call to `make_restrictions`. If you prefer to read good documentation, see the first section of this part of the LISREL documentation. FAiR is attempting to do the same calculations (with some differences in notation), although FAiR does not do unweighted least squares or generalized least squares or multisample models (yet).

The main discrepancy functions used in FAiR are¹

$$\begin{aligned} F_{ML} &= \ln |\mathbf{C}| - \ln |\mathbf{S}| + \text{tr}(\mathbf{S}\mathbf{C}^{-1}) - n, \\ F_{ADF} &= (\mathbf{s} - \mathbf{c})' \mathbf{\Gamma}^{-1} (\mathbf{s} - \mathbf{c}), \end{aligned}$$

The first, F_{ML} , is the maximum likelihood discrepancy function that is familiar in the literature and is used exclusively by the previously-existing R functions for factor analysis `factanal` and by `sem`. One of the sources of added value in FAiR is that users are not limited to F_{ML} which presumes that the data are multivariate normal. Another reason to prefer F_{ML} is that it is a scale invariant function in the case of EFA and in the case of SEFA and CFA if the only exact restrictions are exclusion restrictions. However, as was stated in the Notation section, FAiR utilizes an “embedded correlation” parameterization, i.e. $\mathbf{C} = \mathbf{\Omega}(\beta\Phi\beta' + \Theta^2)\mathbf{\Omega}$, which implies that β and Θ^2 will take the same values regardless of the scale on which the outcome variables are measured.

The second, F_{ADF} , is the “asymptotically distribution free” discrepancy function and is also relatively well-known in the literature (see Browne 1982 and Browne 1984), although not previously available in R. There are a variety of ways to calculate $\mathbf{\Gamma}$, depending on the assumptions one wants to make about the data. Several of which are discussed in Bentler and Dudgeon (1996) and are implemented in FAiR. However, F_{ADF} requires an estimate of $\mathbf{\Gamma}$, which is the subject of the next subsection.

2.1 Estimating Gamma

The process by which $\mathbf{\Gamma}$ is estimated is somewhat convoluted in the source code because $\mathbf{\Gamma}$ is potentially estimated in `make_manifest` and then reestimated in `make_restrictions` when the discrepancy function is chosen. If the raw data are available, $\mathbf{\Gamma}$ is (preliminarily) estimated by `make_manifest`, because it is useful in calculating a robust estimate of the variance-covariance matrix of the parameters, even if F_{ML} will have been chosen as the discrepancy function.

In general, assuming the eighth-order moments of the data are finite

$$\Gamma_{ij,kl} = \sigma_{ijkl} - \sigma_{ij}\sigma_{kl},$$

where σ_{ijkl} is a fourth-order central moment. By default, $\mathbf{\Gamma}$ is calculated with a (biased) maximum likelihood estimator. Let \mathbf{y}_t be a row vector of n observed manifest variables for the t th unit of observation that are deviated from the sample means of the n variables. Let $\mathbf{z}_t = \text{vecs}(\mathbf{y}_t'\mathbf{y}_t)$ be a vector of length n^* . Then, \mathbf{s} is a vector of column means of \mathbf{Z} , where \mathbf{Z} is a matrix that stacks these N vectors, and $\hat{\mathbf{\Gamma}} = \text{cov}(\mathbf{Z})$, where N is used as the divisor instead of $N - 1$. This formula is equivalent to that in Bentler and Dudgeon (1996), except $N - 1$ is used in calculating \mathbf{s} there.

Since $\mathbf{\Gamma} = \text{acov}(\mathbf{s})$, another way to estimate $\mathbf{\Gamma}$ is to simply to bootstrap the estimator of \mathbf{s} . In particular, bootstrapping should probably be undertaken if \mathbf{s} is estimated in some exotic fashion. If the raw data are available, the `bootstrap` argument to `make_manifest` can be specified as some large positive number of bootstraps. Otherwise, the previous estimator of $\mathbf{\Gamma}$ will be executed, if possible. One instance where this estimator is impossible is when some of the outcome variables are ordinal and polychoric and / or polyserial correlations are calculated. In that case, the estimate

¹There is another objective function based on the one in Yates (1987) that is not formally a discrepancy function. It probably should not be used without further refinement.

of $\mathbf{\Gamma}$ is diagonal, with the diagonal consisting of the squared standard errors of each element of \mathbf{s} . This discrepancy function is sometimes known as “diagonally-weighted least squares” (DWLS).

However, if the raw data are available and all variables are continuous, this estimate of $\mathbf{\Gamma}$ can be superceded in the call to `make_restrictions` when the discrepancy function is "ELLIPTICAL", "HK", or "SHK". In the heterogenous kurtosis (HK) case

$$\begin{aligned}\sigma_{ij,kl} &= (a_{ij}a_{kl})\sigma_{ij}\sigma_{kl} + (a_{ik}a_{jl})\sigma_{ik}\sigma_{jl} + (a_{il}a_{jk})\sigma_{il}\sigma_{jk}, \\ a_{ij} &= \frac{\kappa_i + \kappa_j}{2}, \\ \kappa_i &= \sqrt{\frac{\sum_{t=1}^N (y_{ti})^4}{3 \sum_{t=1}^N (y_{ti})^2}}.\end{aligned}$$

Note that κ_i is a kurtosis estimate. If $\kappa_i = \kappa \forall i$, then the elliptical case results, and if $\kappa_i = 1 \forall i$, the multivariate normal case results. FAiR includes a “shrunk heterogenous kurtosis” (SHK) estimator that shrinks each κ_i towards the median kurtosis estimate. However, to explain the shrinking process, you have to read the next section.

3 Using `make_manifest()`

To estimate a factor analysis model, it is first necessary to obtain an estimate of $\mathbf{\Sigma}$ from the data, denoted \mathbf{S} . The `make_manifest()` generic function does so and has three primary, and somewhat redundant arguments, `x`, `data`, and `covmat`. The following table may clear things up somewhat:

x	data	covmat	comment
missing	missing	list	must have <code>cov</code> element
missing	missing	hetcor	probably better to pass the data instead
missing	missing	CovMcd	probably better to pass the data instead
missing	missing	matrix	should specify <code>n.obs</code> also
matrix, data.frame	missing	missing	all variables in <code>x</code> will be used
missing	matrix, data.frame	missing	all variables in <code>data</code> will be used
formula	data.frame	missing	formula should not contain a response

The methods above the double line are used when `x` and `data` are unspecified but `covmat` is specified. These methods pertain to the situation in which a covariance “matrix” (in some form) is passed to `make_manifest()` via `covmat` but the raw data are not available. The `covmat` argument could be a covariance matrix (or a correlation matrix, in which case it would be preferable to also specify the `sds` argument, which is a vector of manifest standard deviations) or could be an object that contains a covariance matrix, such as a list (with an element named `cov`), an object of S3 class “hetcor” (produced by the suggested polycor package) or an object of S4 class `CovMcd` (produced by the rrcov package). However, it is usually unnecessary to make use of the latter two options because `make_manifest()` can produce such objects if the raw data are passed to it (see below).

If `covmat` is a list, it may also contain an element named `n.obs`, which is the number of observations, and an element named `W`, which is an estimate for $\mathbf{\Gamma}^{-1}$ to be used with F_{ADF} . Again, this mechanism should rarely need to be used because FAiR can estimate $\mathbf{\Gamma}$ in several ways, provided that the raw data are available.

The methods below the double line in the above table are used when the raw data are available and have a few additional arguments that govern how \mathbf{S} is to be calculated, whether bootstrapping should take place, etc. The most common estimator is the unbiased one, which is simply equal to the inner products of pairs of centered variables, divided by $N - 1$. The maximum likelihood estimator is similar but divides by N instead. Either is appropriate under ideal conditions, but such conditions are rare. Hence, `make_manifest()` has several alternative estimators of $\mathbf{\Sigma}$.

Two kinds of “shrinkage” estimators of Σ are implemented in `make_manifest()`. There are two reasons why one might be interested in a shrinkage estimator. First, the unbiased estimator produces biased estimates of the eigenstructure, particularly when $\frac{N}{n}$ is small. Second, if we assume that the number of “common” factors in the data are greater than r but that the excess minor factors tend to exacerbate observed correlations, then it may be reasonable to fit a covariance matrix that has been “shrunk” a bit toward an identity matrix.

The first shrinkage estimator makes use of the `cov.shrink` function in the suggested `corpcor` package; see also SchÄfer and Strimmer (2005) and Opgen-Rhein and Strimmer (2007). Let

$$\begin{aligned}\hat{\lambda}_1^* &= \min \left(1, \frac{\sum_{i \neq j} \widehat{var}(r_{ij})}{\sum_{i \neq j} r_{ij}^2} \right), \\ \hat{\lambda}_2^* &= \min \left(1, \frac{\sum_{i=1}^n \widehat{var}(v_i)}{\sum_{i=1}^n (v_i - v_m)^2} \right),\end{aligned}$$

where r_{ij} is an unbiased sample correlation estimate and v_i is an unbiased sample variance estimate and v_m is the median of these variance estimates. Then,

$$\begin{aligned}r_{ij}^* &= (1 - \hat{\lambda}_1^*) r_{ij}, \\ v_i^* &= \hat{\lambda}_2^* v_m + (1 - \hat{\lambda}_2^*) v_i, \\ s_{ij}^* &= \frac{r_{ij}^*}{\sqrt{v_i^* v_j^*}}.\end{aligned}$$

This estimator of Σ can be obtained by specifying `how = "lambda"` in the call to `make_manifest` when the raw data are available.

The same idea can be applied to obtain a “shrunk heterogenous kurtosis” (SHK) estimator of Γ , which may have a smaller root mean-squared error than HK estimator. Let

$$\begin{aligned}\kappa_i &= \sqrt{\frac{\sum_{t=1}^N (y_{ti})^4}{3 \sum_{t=1}^N (y_{ti})^2}}, \\ \hat{\lambda}_2^* &= \min \left(1, \frac{\sum_{k=1}^n \widehat{var}(\kappa_i)}{\sum_{k=1}^n (\kappa_i - \kappa_m)^2} \right), \\ \kappa_i^* &= \hat{\lambda}_2^* \kappa_m + (1 - \hat{\lambda}_2^*) \kappa_i,\end{aligned}$$

and $a_{ij}^* = \frac{\kappa_i^* + \kappa_j^*}{2}$ is used — rather than a_{ij} — to estimate of Γ in the last formula in the previous section. If $\hat{\lambda}_2^* = 1$, then something quite similar to the elliptical estimator would result, so FAiR simply reverts to the elliptical estimator in that case.

The second “shrinkage” estimator is simpler and can be applied in a post-hoc fashion to (almost) any \mathbf{S} by specifying `shrink = TRUE` in the call to `make_manifest`. The second shrinkage estimator preserves the eigenvectors of \mathbf{S} but compresses the eigenvalues, which is proven to be a minimax estimator under the conditions specified in theorem 3.1 of Dey and Srinivasan (1985). Let $\lambda(\mathbf{S})$ be the eigenvalues of the preliminary \mathbf{S} , arranged in descending order. The j th eigenvalue of the final \mathbf{S} is set equal to $\frac{N\lambda_j(\mathbf{S})}{N+n+1-2j}$ and then the final \mathbf{S} is reassembled by multiplying the diagonal matrix of new eigenvalues from the left and right by the old eigenvector matrices. This shrinkage estimator can often be used even when the raw data are not available, and the preliminary \mathbf{S} was passed to `covmat`.

The “minimum covariance determinant” (MCD) estimator is probably the best choice when all variables are continuous, particularly if there may be multivariate outliers in the data. This estimator can be obtained by specifying `how = "MCD"` in the call to `make_manifest`. In short, the MCD estimator picks a subsample of observations such

that the determinant of the preliminary \mathbf{S} is minimal. Then, a variety of correction factors are applied to dramatically improve the efficiency of the estimator without undermining its high break-down point. The details are well explained in Rousseeuw and van Driessen (1999), in Pison, Van Aelst, and Willems (2002), and in the documentation of the `CovMcd` function in the `rrcov` package.

Another alternative when there are outliers is to fit a matrix of Spearman correlations, which can be obtained by specifying `how = "ranks"` in the call to `make_manifest`. This ranks-based estimator has another important role; namely it is used when bootstrapping a polychoric correlation matrix. In that case, the polychoric correlations are obtained using the `hetcor` function in the suggested `polycor` package. Then, a large number of Spearman correlation matrices are obtained via bootstrapping, and finally the bias in the bootstrap distribution is corrected by rescaling the distribution so that its means are equal to the polychoric point estimates. This short-cut procedure is reasonably effective and considerably faster than bootstrapping a polychoric correlation matrix when there are a large number of observations, a large number of variables, and a large number of categories for each variable.

If all the variables are continuous but there are missing data, then the `mlest` function in the suggested `mvnmle` package is used to estimate Σ assuming multivariate normal data and Missing At Random (MAR) missingness. See its documentation for more details.

For all these estimators of Σ — except the unbiased and maximum likelihood estimators — we do not know the asymptotic covariance of \mathbf{s} and such an estimate is necessary to use the ADF discrepancy function. Thus, it is best to bootstrap the estimate of Γ in those cases, which can be done by setting `bootstrap` equal to the number of desired bootstraps in the call to `make_manifest`. One can also specify `wt`, which governs the probability that an observation will be selected in the nonparametric bootstrap sampling. The `wt` argument can also be used with the maximum likelihood or unbiased estimators of Σ when the sample has probability weights.

4 Esoteric definitions

Aside from the discrepancy functions and test statistics, FAiR differs from the implementation described in other software manuals, textbooks, etc. Thus, it is important to highlight three concepts that are perhaps not well-known among factor analysts.

4.1 The factor contribution matrix

As stated in the Notation section, the factor contribution matrix is defined (see White 1966 and Yates 1987) as the element-by-element product of a pattern and a structure matrix, i.e. $\mathcal{D} = \beta \times \Pi$, where $\Pi = \beta\Phi$, and similarly for level two. Hence if $\beta_{jp} = 0$, then $\mathcal{D}_{jp} = 0$, but $\Pi_{jp} = 0$ also implies $\mathcal{D}_{jp} = 0$.

Each row of \mathcal{D} sums to the communality of the corresponding outcome variable. As White (1966) notes, each cell of \mathcal{D} “represents the proportion of variance on test j associated with variation on factor p . Specifically, it is the decrement of [variance in the j th outcome variable] which would occur if the [p th factor were] dropped from the specification ... without adjusting the factor loadings on the other factors. (375)”

The factor contribution matrix was originally derived because some thought it would aid in the interpretation of factors. The interpretation of \mathcal{D}_{jp} is straightforward, since it is just the correlation between the j th outcome and the p th factor, weighted by β_{jp} . Also, the column-wise sums of \mathcal{D} provide a measure of the “strength” of the p th factor in the battery; in fact, by default the factors produced by `summary(*)` are presented in decreasing order of the column-wise sums of \mathcal{D} . This matrix of factor contributions can be extracted for interpretation using `loadings(*, matrix = "FC")` but has additional roles in FAiR.

If $r > 1$ and $\Phi \neq \mathbf{I}$, then any cell of \mathcal{D} can be negative, which as Yates (1987) notes, implies that the p th factor is a negative suppressor variable for the j th outcome and has negative weighted validity (Conger 1974). Conversely, we could say that outcome j is the “best” indicator (in the sense of having the most weighted validity) of factor p if $\mathcal{D}_{jp} = \max(\mathcal{D}_p)$.

As discussed below, many restrictions that can be imposed on a factor analysis model are defined with respect to \mathcal{D} in FAiR, rather than with respect to β . Also, following a suggestion emailed to me by Peter Bentler, it is possible to use \mathcal{D} rather than β as the relevant matrix for some analytic transformation criteria for EFA.

4.2 Generalized variance (GV) and effective variance (EV)

The generalized variance of a set of variables is a generalization of the univariate concept of variance and is defined as the determinant of their covariance (or correlation) matrix. The GV can be interpreted as the hypervolume of the variables. In FAiR, the relevant matrices are all correlation matrices, and hence the GV is defined here on standardized variables. For example, $|\Phi|$ is the GV of the primary factors.

One limitation of the GV is that it is inappropriate for comparisons between sets of variables that are of different sizes. Hence, Peña and Rodríguez (2003) derives the “effective variance” of a set of variables as the determinant of their covariance matrix raised to the power of the reciprocal of their number. For example, $|\Phi|^{\frac{1}{r}}$ is the EV of the primary factors. Peña and Rodríguez (2003) suggests that the EV can be interpreted as the edge-length of a hypercube whose volume is the GV of the variables and shows that it is a valid metric for comparing dispersion across differently-sized sets of variables.

I have never encountered a paper in the factor analysis literature that attempted to interpret a result on the basis of the GV or EV, although it is in harmony with the geometric interpretation of the factor analysis model. Nevertheless, both have important roles in FAiR because restrictions can be defined on them, particularly in the case of the EV. Yates (1987) advocates one such restriction, and several more have been included in FAiR that are loosely derived from Yates (1987) are discussed below.

5 Important theorems

The two (three really) theorems discussed in this section are relevant to the identification of CFA and SEFA models.

5.1 Howe (1955) and Koopmans and Reiersøl (1950)

One theorem from Howe (1955) is somewhat well-known via various works of Karl Jöreskog. A set of conditions that is sufficient for rotational uniqueness of β are that

1. Φ is a full-rank correlation matrix
2. β has $r - 1$ zeros in each column
3. β^p is of rank $r - 1$, where β^p is the submatrix of β that includes all rows where the p th factor is zero

Bollen and Jöreskog (1985) and Millsap (2001) emphasize that these conditions are insufficient for identification of the parameters. Rotational uniqueness roughly means that there is no transformation matrix that can post-multiply β , leave the zeros in the same positions, and change the values of the nonzeros. Identification roughly means that one could uniquely solve for all free model parameters if the population Σ were available.

A similar theorem was proven in Koopmans and Reiersøl (1950), although this paper has been more influential in the econometrics literature than in the factor analysis literature. I think Koopmans and Reiersøl (1950) should be credited with Howe’s (1955) result, although it is not clear if Howe (1955) was aware of it. Koopmans and Reiersøl (1950) shows that if Θ^2 is identified and the $r - 1$ zeros in β are specified in specific cells *a priori*, then these conditions are collectively necessary and sufficient for identification of the free parameters in β and Φ .

In the example from Bollen and Jöreskog (1985), Θ^2 is not identified, and thus neither are β and Φ . If $r \leq 2$ or $r = 3$ and $n \geq 7$, then a necessary and sufficient condition for the identification of Θ^2 is that, after deleting any row of β , there remain two disjoint submatrices of rank r , which is a well-known result from Anderson and Rubin (1956). If $r > 3$, this condition is merely sufficient, but it is a rather weak sufficient condition.

The upshot of Koopmans and Reiersøl's (1950) result is that it gives minimal conditions under which a CFA model is identified. Warnings will be produced if a CFA model appears to be inconsistent with these conditions. Note, however, that if $r > 2$, it is not possible to verify the third condition with certainty. FAiR will check that the rank of each β^p could be $r - 1$ but cannot preclude the possibility that two or more columns of β^p besides column p are linear functions of each other in the population.

5.2 Reiersøl (1950)

Koopmans and Reiersøl (1950) also states another important theorem that is more relevant for SEFA, although the proofs are given in Reiersøl (1950).

1. Φ is a full-rank correlation matrix
2. β has r zeros somewhere in each column (without committing to where they are located)
3. β^p is of rank $r - 1 \forall p$, where β^p is the submatrix of β that includes all rows where the p th factor is zero
4. If any row of β that is not in β^p is appended to β^p , the rank of the resulting matrix is r
5. β_{-t}^p is of rank $r - 1 \forall p, t$, where β_{-t}^p is the submatrix of β^p that excludes the t th row
6. No other $r \times r$ submatrix of β is of rank $r - 1$ besides the $\beta^1 \dots \beta^r$ matrices
7. Θ^2 is identified

Reiersøl (1950) contains a weaker theorem, but it is characterized as “less useful” and is not considered here (yet). It is possible for a SEFA model to enforce conditions 1 and 2 by construction. FAiR will check that the ranks of β^p and β_{-t}^p could be $r - 1$ but again cannot preclude the possibility that the rank is actually smaller in the population. Similarly, it is impossible to verify conditions 4 and 6 without population knowledge, but we usually are not too concerned with such corner cases.

The upshot of Reiersøl's (1950) theorem is that it provides minimal (albeit extensive) conditions for identification in SEFA models where the number of zeros in the p th column of β is specified but not their locations. Hence, the prototype value for the `rankcheck` slot of an object that inherits from class `paramter.coef.SEFA` is "reiersol" for SEFA models and warnings will be produced if the model appears to be unidentified. It is possible that identification could plausibly be asserted under various weaker conditions, particularly if inequality restrictions are used to preclude some otherwise duplicative solutions. However, departures from Reiersøl's (1950) theorem would have to be considered on a case-by-case basis.

6 SEFA or CFA via `Factanal()`

In this example, I will use a covariance matrix containing 24 mental tests used by Harman (1976). Execute `?Harman74.cor` to see more information about it, but it is just a fairly standard and well-known mental test example.


WARNING: The editable menus are kept small in this vignette; it may be necessary to resize the editor window or scroll to access all the cells of the matrix you intend to edit. Also, note that changes to a cell do not “register” until you press enter, tab, arrow, etc. If you have an “unregistered” change and click OK, that change will be ignored. Also, if you click Cancel, the function aborts rather than reverts to the previous menu.

To start this sequence, I typed the commands

```
library(FAiR)
man <- make_manifest(covmat = Harman74.cor)
res <- make_restrictions(manifest = man, model = "SEFA")
```

6.1 Number of Factors

The first questions that will be asked pertain to the number of factors and can be avoided if factors were specified explicitly, e.g. `make_restrictions(manifest = man, factors = c(5,2), model = "SEFA")`.




Some people may be unfamiliar with the idea of a two-level model; it is briefly defined in the Notation section and also in the help page for `make_restrictions`. If $r \geq 3$, the user will be asked whether to estimate a simultaneous second-order model, which decomposes the correlation matrix among first-order primary factors (Φ) as a function of fewer second-order factors. For example, some people define “general intelligence” to be the second-order factor that drives various first-order (primary) mental abilities.

If a second-order model is estimated, and if $r \geq 5$, the user will be asked how many second-order factors (\tilde{r}) to extract. For the sake of completeness, I will assume that the user answered “Yes” to the question of whether a simultaneous second-order model should be estimated. Then, the dialog asks the user to specify the number of second-order factors to extract. If $r = 3$ or $r = 4$, then \tilde{r} can only be one. If $r = 5$, then \tilde{r} can be one or two. If $r \geq 7$, then \tilde{r} can be three in a SEFA model. Rarely would one need to estimate a model with larger numbers for r and \tilde{r} . I will assume that the user has chosen $\tilde{r} = 2$.


6.2 Main menu

The following menu is critical in the sense that the user will only be subsequently asked about options that are checked. This menu would have fewer options if there were no simultaneous second-order model, or if there were only one second-order factor, or if the model were a CFA. It is quite possible to check multiple options. I will discuss each of these options eventually; however, the order in which subsequent options are specified is not necessarily the same order that is presented in this menu.



6.2.1 Fix specific coefficients to particular values (second and eighth checkboxes)

The figures below illustrate dialogs that ask about pegging specific coefficients to particular values *a priori* and hence not estimating them. The dialogs for both level two ($r \times \tilde{r}$) and level one ($n \times r$) are shown simultaneously here but are not actually simultaneously specified when using FAiR. Although it is not visibly apparent unless the editor window is maximized, the dimensions of matrix to be edited are the same as the dimensions of the matrix to eventually be estimated, and there is an exact correspondence between the cells of the editor window and the cells of β (or $\tilde{\beta}$ as the case may be). This correspondence holds in almost all of the editor windows to follow, but this matrix-based interface may not be as intuitive to users who are accustomed to specifying SEMs with path diagrams. Although this example is a SEFA, these dialogs are more characteristic of a CFA model, which requires enough *a priori* zero coefficients to satisfy the theorems discussed in section 5.1. If this were a CFA model, these editing windows would pop up regardless of whether the user requested so.



There are several important things to note. First, any unrestricted coefficients should be left as NA. In many SEM programs, a coefficient is restricted to be zero unless the user specifies otherwise. This is *not* the case in FAiR; if and only if you specify that a coefficient is zero will it be zero *a priori*. In a SEFA model, it is possible to leave all cells as NA, but doing so would generate an error message under a CFA model.

Second, in many SEM programs it is possible or even customary to specify that some coefficient is 1.0 to normalize the scale of a factor. In FAiR, the user generally should *not* restrict one coefficient per factor to have a coefficient of 1.0 because the factors are already normalized to have unit variances along the diagonal of Φ . Hence, to also constrain a coefficient to be 1.0 would be a substantive restriction rather than merely a normalization.

Third, if one were to restrict a coefficient to be 1.0, then one would almost always restrict the other coefficients in that row to be 0.0, in which case the corresponding outcome variable will be treated as “exogenous” and its standard deviation will be estimated from the data rather than by the model. Hence, this mechanism (which is apparently all too familiar to long-time users of LISREL) can be used for factor analysis where some factors are identical to observed manifest variables. However, this specification is not generally recommended in FAiR. It is better to exploit a feature of FAiR (discussed below) to designate some outcome variable as the “best” (yet imperfect) indicator of a factor.

Fourth, if one wanted to impose the restriction that some coefficient is an exact function of other coefficients (but not the simpler restriction that two or more coefficients are equal), then it is necessary to designate the “fixed” coefficient in this dialog and pass a function that enforces these restrictions to the `nl_1` or `nl_2` arguments of `make_restrictions`. For example, to impose the restriction that the first coefficient on the “Visual Perception” test is the ratio of the second and third coefficients, one could proceed like this

```
foo <- function(x) {
  x[1,1] <- x[1,2] / x[1,3]
  return(x)
}
res <- make_restrictions(manifest = man, model = "SEFA", nl_1 = foo)
```


But it is also necessary to designate β_{11} as “fixed”, even though we do not know what number it is fixed to until β_{12} and β_{13} are specified. In such situations, “fix” coefficients that are exact functions of other coefficients to `Inf` (note the capitalization of `Inf`).

Fifth, if merely imposing the restriction that two or more coefficients are equal to each other, do not pass such a function to `nl_1` or `nl_2` and do not bother designating affected coefficients as “fixed” in this dialog. Instead, use the dedicated mechanism for imposing equality restrictions, which is discussed in the next subsection.

All that said, the most common use of this dialog (particularly for a CFA) is to restrict specific coefficients to be 0.0. Simply type 0.0 in the appropriate cell(s), *press Enter to make sure the change “registers” in the editor*, and press OK to exit this dialog. If restricting a specific coefficient to some number other than 0.0, keep in mind that the embedded correlation parameterization in FAiR means that coefficients are calibrated for *standardized* variables.

6.2.2 Constrain coefficients to be equal to each other (fourth checkbox)

This figure illustrates the dialog to constrain one or more coefficients (at level one) to be equal to each other; the dialog for level two is similar. Again, it is not visibly apparent unless the editor window is maximized but the dimensions of the matrix to be edited are $n \times r$ just like β .



The matrix is initially filled with FALSE, which is a keyword in R. To impose an equality restriction among two or more coefficients, change at least two of the cells to TRUE, which is also a keyword in R. *The capitalization is essential.* For example, to impose the restriction that $\beta_{11} = \beta_{21}$, I would change the first two cells in the first column to TRUE. It is possible to constrain three or more coefficients to all be equal to each other by changing three or more cells to TRUE. When all cells that are restricted to be equal to each other are designated with TRUE in the appropriate cell, click OK.

At that point, all the cells will be reset to FALSE, which allows you to impose *another* set of equality restrictions (among a disjoint set of coefficients, e.g. $\beta_{14} = \beta_{15}$). This process can be repeated as many times as is necessary to represent all the equality coefficients you want to impose on disjoint sets of coefficients. If OK is clicked when all cells remain FALSE, then whatever dialogue box is next appears.

Finally, again recall that the embedded correlation parameterization in FAiR implies that coefficients should be calibrated for standardized variables. Hence, constraining two or more cells of β to be equal does not imply that the corresponding cells of $\Omega\beta$ will be equal.

6.2.3 Change bounds on coefficients (third checkbox)

These figures ask about bounds on coefficients at level one, i.e. on β . The dialogue for level two would be similar.

Please select (multiple selections possible)

Are you sure you would like to place non-trivial bounds on some of the (free) coefficients at level 1 ?

☒ Change lower bound (currently -1.5)

☐ Change upper bound (currently 1.5)

OK Cancel

Please select one

What number would you like to use as a lower bound for **most** of the coefficients at level 1 ?

-0.1

OK Cancel

Editing needed

Please edit this matrix to specify lower bounds on each coefficient at level 1. Recall that coefficients are parameterized for **standardized** variables. Press OK when finished.

Row.names	F1	F2	F3	F4	F5
VisualPerception	-0.100000	-0.100000	-0.100000	-0.100000	-0.100000
Cubes	-0.100000	-0.100000	-0.100000	-0.100000	-0.100000

OK Cancel

Specifying bounds on coefficients is a three-stage process. First, the user is asked whether to change the lower and / or upper bounds from their default values, here ± 1.5 . I only illustrate changing the lower bounds, but the process is similar if one wanted to additionally or alternatively change the upper bounds.

Second, the user is asked for a “fill value” to use for *most* of the (lower in this case) bounds on coefficients. After pressing OK, this value is filled into an editable matrix that is of the same dimension as β , although that is not obvious when the editor window is not maximized. Third, the user can edit any cell of this $n \times r$ matrix to modify the bound for any coefficient. Here I leave all lower bounds at my “fill value” of -0.1 . Thus, this model requires a “almost-positive” manifold, which is a *substantive* constraint that can be wrong and can be tested.


Finally, again recall that the embedded correlation parameterization in FAiR implies that coefficients (and hence their bounds) should be calibrated for standardized variables. In particular, rarely would it be necessary to make the bounds *wider* than ± 1.5 because more extreme standardized coefficients are incredibly rare (provided that Φ is well-conditioned). Moreover, if there were only one factor, it is impossible to make the bounds on its coefficients wider than ± 1.0 . If one were to impose the inequality restriction that there are no negative suppressor variables (see below), then it is somewhat advantageous to also narrow the upper bound on the coefficients to 1.0.

6.2.4 Important Note

The running example is a two-level model. In two-level models, one cannot currently impose bounds on the off-diagonals of Φ using the pop-up menus. However, if this were a one-level model, the next dialog box would ask about bounds on the off-diagonals of Φ if such an option were selected from the main menu. That dialog would be similar to the one illustrated in subsection 6.2.6.

6.2.5 Nondefault mapping rule and changing the requisite number of zeros (11th and 12th checkboxes)

In the SEFA model presumed here, the question shown in the figure below is very important and pertains to the mapping rule used to squash certain cells of β to zero during the optimization. The dialog for level two is similar but generally has fewer options. If a CFA model were being estimated, this question would not appear because the theorems in section 5.1 must be satisfied by *a priori* restrictions. Specifying a non-default mapping rule is more common for a rather “exploratory” SEFA; for a more “confirmatory” SEFA, one would typically accept the default mapping rule and supplement it with other restrictions.



Let b_p be the number of zeros required for the p th column of β . Let $\underline{\beta}$ be a matrix with the same dimensions as β . Suppose the free cells of $\underline{\beta}$ have been filled in by a proposal for the restricted optimum of the discrepancy function, but it has fewer than b_p zeros in the p th column (and possibly no zeros at all). The “mapping rule” maps from $\underline{\beta}$ to β by changing some “small” cells of $\underline{\beta}$ to zero and then the discrepancy function (and each constraint function) is evaluated at β , not $\underline{\beta}$. By “small”, I always mean in absolute value; however the rules are usually defined on the reference structure matrix ($\underline{\Upsilon} = \underline{\beta}\underline{D}$) or the factor contribution matrix ($\underline{\vartheta} = \underline{\beta} \times (\underline{\beta}\underline{\Phi})$), rather than $\underline{\beta}$ itself.

- “No, I just want plain zeros”. This default mapping rule simply squashes the b_p smallest elements in $\underline{\beta}_p$ to zero.
- “At least one zero per factor on a high communality variable.” This mapping rule is similar but is slightly more complicated. Let $\overline{\vartheta}_j = \frac{1}{r} \sum_{p=1}^r \underline{\vartheta}_{jp} = \frac{h^2}{r}$ be the arithmetic mean of the factor contributions for the j th outcome variable and let $\widetilde{\vartheta}_j = \left(\prod_{p=1}^r \underline{\vartheta}_{jp} \right)^{\frac{1}{r}}$ be the geometric mean of the factor contributions for the j th outcome, which is defined only for outcomes with nonnegative factor contributions. Hence, $\frac{\overline{\vartheta}_j}{\widetilde{\vartheta}_j}$ will tend to be more positive when the communality of the j th outcome is large but at least one factor contributes little to the

communality. Define the set of “potential zeros” on factor p as those rows of $\underline{\mathbf{Y}}$ where $\underline{Y}_{jp} < \underline{Y}_{jq} \forall q \neq p$. For each p , this mapping rule squashes $\underline{\beta}_{jp}$ to zero for the outcome among the set of “potential zeros” with the largest (well-defined) $\frac{\underline{\partial}_j}{\underline{\partial}_p}$. The default mapping rule is then applied to squash $b_p - 1$ more cells to zero for the p th factor.

- “Encourage cohyperplanarity (revisionist Yates 1987)”. This very promising mapping rule squashes $r - 1$ cells in the p th column of $\underline{\beta}$ using the following piecewise function $\forall q \neq p$:

$$\beta_{jp} = \begin{cases} 0 & \text{if } \underline{\partial}_{jq} - \underline{\partial}_{jp} = \max \{ \underline{\partial}_q - \underline{\partial}_p \} \\ \underline{\beta}_{jp} & \text{otherwise,} \end{cases}$$


which tends to place the zeros for the p th column of $\underline{\beta}$ in the rows where the factor contribution of the q th factor is large. The default mapping rule is then applied to squash $b_p - (r - 1)$ more cells to zero for the p th factor. Although Yates (1987) did not anticipate SEFA, it repeatedly recommends procedures that tend to encourage outcomes within hyperplanes in the common factor space to be widely dispersed throughout their hyperplane(s). This mapping rule attempts to do so.

- “ r outcomes of complexity $r - 1$ for each factor (weaker Thurstone)”. This mapping rule is similar to the default mapping rule in that r small elements in the p th column of $\underline{\beta}$ are squashed to zero, except that there is an additional stipulation that no outcome is of complexity less than $r - 1$ where the complexity of the j th outcome is its number of non-zero coefficients. However, any “non-zero” coefficient can be arbitrarily close to zero. Thus, each column of $\underline{\beta}$ will have r *exact* zeros. If necessary, the default mapping rule will then be applied to squash $b_p - r$ more cells to zero for the p th factor.
- “ $2r$ outcomes each of complexity $r - 2$.” This mapping starts by finding the smallest cell in the first column of $\underline{\mathbf{Y}}$ and squashes it to zero. Say this cell is \underline{Y}_{j1} . Then, the smallest nonzero cell in the j th row of $\underline{\mathbf{Y}}$ is also squashed to zero so that the j th outcome is of complexity $r - 2$. This procedure is repeated for all columns to yield $2r$ outcomes each of complexity $r - 2$, such that each factor has $r - 1$ zeros. Finally, the default mapping rule is applied to squash $b_p - (r - 1)$ more cells to zero for the p th factor.
- “Unit complexity basis (revisionist Butler 1969)”. Butler (1969) defines “the r most distinguished outcomes” and proposes to run the r primary axes through them in EFA to create a unit complexity basis for $\underline{\beta}$. The following SEFA mapping rule is similar and is defined by a piecewise function $\forall q \neq p$:


$$\beta_{jp} = \begin{cases} 0 & \text{if } \text{abs}(\underline{\beta}_{jq}) = \max(\text{abs}(\underline{\beta}_q)) \\ \underline{\beta}_{jp} & \text{otherwise} \end{cases}$$

The default mapping rule is then applied to squash $b_p - (r - 1)$ more cells to zero for the p th factor.

- “Set the maximum complexity for each outcome.” This mapping rule is similar to the default mapping rule, except that it is applied to the *rows* of $\underline{\mathbf{Y}}$ rather than its columns. In general, $\forall j$, the user can choose $b_j \in \{0, \dots, r - 1\}$ for the minimum number of zero coefficients in $\underline{\beta}_j$, and the complexity of the j th outcome is $r - b_j$. It is possible to choose the same complexity for all rows of $\underline{\beta}$. If the complexity of $\underline{\beta}_j$ is $r - 1 \forall j$, this restriction characterizes the main tenet of Thurstone’s (1935) definition of simple structure. If the complexity of $\underline{\beta}_j$ is $1 \forall j$, this restriction implies a perfect cluster configuration. Alternatively, b_j can be specified on a row-by-row basis, as is shown in the dialog below, which will pop up if this mapping rule is selected. Finally, the default mapping rule is applied to ensure that there are at least b_p zeros for the p th factor.




Finally, in the case of SEFA, it is necessary to specify $b_p \forall p$. By default, $b_p = r \forall p$, which is the minimum that is necessary for identification of the nonzero parameters. However, it is possible to select $b_p > r$ and may be possible to select $b_p = r - 1$ in some cases if identification of the relevant parameters can be accomplished in some other fashion. The following dialogs illustrate this mechanism, which is initialized when the box for “Change the number of zero coefficients required for each factor” is checked in the main menu:



6.2.6 Change bounds on correlations among primary factors (first checkbox)

This figure below illustrates a dialog that asks about bounds on the off-diagonals of $\tilde{\Phi}$. If $\tilde{r} \leq 1$, this question would not be asked, although if $\tilde{r} = 0$, a similar dialogue would ask about bounds on the off-diagonals of Φ . If there were more factors, the editor window would have more rows, totalling $0.5\tilde{r}(\tilde{r} - 1)$.

The user is given an editor with columns for the lower and upper bounds on these correlations. The bounds cannot exceed ± 1 because they pertain to correlations. Here, I edited the admissible interval to $[-0.9, 0.9]$, which is a largely irrelevant restriction.




6.2.7 Impose functional inequality restrictions

It is always important to keep in mind that FAiR does something tantamount to restricted optimization and that the possibilities for restrictions are endless. Hence, perhaps the most important feature of FAiR is the ability to impose inequality restrictions on functions of multiple parameters. It is even possible to define your own restriction functions and pass them to the criteria argument of `make_restrictions`. In this subsection, I will only discuss the functional inequality restrictions that are pre-programmed in FAiR.

Please refer to section 4 for definitions of the factor contribution matrix, the generalized variance, and the effective variance. Also, let $\mathbb{I}\{\cdot\}$ be the “indicator function”, which equals 1 if the condition inside the braces is true and equals 0 if the condition is false. In FAiR, an inequality restriction function is a piecewise function that equals -1 if the restriction is satisfied and equals some number greater than -1 if the restriction is not satisfied. You do not need to worry too much about the return values for the case where the restriction is not satisfied.

When the box for “Impose functional inequality restrictions” (at level one) is checked, the following menu appears:




The menu would be similar at level two but will perhaps have fewer options. These restrictions are defined as follows:

- “Factor contributions must respect row-wise order constraints.” Let \mathbf{G} be a $n \times r$ matrix that is specified by the user to indicate the required ranks of the factor contributions within each row with larger factor contributions receiving smaller ranks. For example, \mathbf{G}_j could be $\begin{bmatrix} 4 & 1 & 5 & 3 & 2 \end{bmatrix}$ to require that the second factor contribution in the j th row be biggest, the third factor contribution be smallest, etc. However, it is not necessary for the ranks to be unique. For example, \mathbf{G}_j could be $\begin{bmatrix} 5 & 1 & 5 & 5 & 5 \end{bmatrix}$ to require that the second factor contribution in the j th row be biggest, while allowing the other ranks to be unrestricted. Such a specification would often be used in CFA or in a more “confirmatory” SEFA where the user has strong theoretical reason to believe that the j th outcome largely “measures” the p th factor and thus specifies that $G_{jp} = 1$ and that $G_{jq} = r \forall q \neq p$. But keep in mind that, for example, $\mathbf{G}_j = \begin{bmatrix} 2 & 2 & 5 & 5 & 5 \end{bmatrix}$ is also valid and requires that the first two factor contributions in the j th row be the top two without taking a position on whether the first is larger than the second. Conversely, $\mathbf{G}_j = \begin{bmatrix} 2 & 2 & 3 & 3 & 5 \end{bmatrix}$, for example, is invalid because it is impossible for the first four factor contributions to all be ranked in the top three.

\mathbf{G} can be specified interactively in response to the following dialog:²

²This restriction can also be enforced from the command line with `make_restrictions(*, criteria = list("ranks_rows_1st"), methodArgs = list(row_ranks_1st = G))` where \mathbf{G} matrix is manually defined in the global environment.




Let $\underline{\mathcal{Q}}$ be a factor contribution matrix that has been converted to ranks within rows. The function that imposes this constraint is formally defined as

$$\frac{-1}{nr} \sum_{j=1}^n \sum_{p=1}^r \mathbb{I} \left\{ \underline{\mathcal{Q}}_{jp} \leq G_{jp} \right\} \in [-1, 0]$$

and thus equals -1 if and only if $\underline{\mathcal{Q}}$ is consistent with \mathbf{G} in every cell.

- “Factor contributions must respect column-wise order constraints.” Essentially, this constraint involves the same concept as the previous constraint, except that $\underline{\mathcal{Q}}$ and \mathbf{G} are defined on the basis of column-wise ranks. I use the same notation as before but the matrices have different contents. For example, $\mathbf{G}'_p = [3 \ 3 \ 3 \ n \ \dots \ n]$ would require that the first three outcomes have the three largest factor contributions on the p th factor. The remaining $n - 3$ outcomes can be ranked in any fashion on the p th factor, provided that none is in the top three. Again, this mechanism would most often be used in CFA or in a more “confirmatory” SEFA. It is tempting to specify that $G_{jp} = 1$ in order to make the p th factor closely related the to j th outcome, but there is a better mechanism for doing so, which is discussed next.
- \mathbf{G} can be specified interactively in response to the following dialog:³

³This restriction can also be enforced from the command line with `make_restrictions(*, criteria = list("ranks_cols_1st"), methodArgs = list(col_ranks_1st = G))` where \mathbf{G} matrix is manually defined in the global environment.



Let $\underline{\vartheta}$ be a factor contribution matrix that has been converted to ranks within columns. The function that imposes this constraint is formally defined as


$$\frac{-1}{nr} \sum_{j=1}^n \sum_{p=1}^r \mathbb{I} \left\{ \underline{\vartheta}_{jp} \leq G_{jp} \right\} \in [-1, 0]$$

and thus equals -1 if and only if $\underline{\vartheta}$ is consistent with \mathbf{G} in every cell.

- “Designate which outcome variable is the best indicator of each factor.” This criterion just is a special case of the previous one but is operationalized differently for better computational performance. In essence, one is merely requiring that $G_{jp} = 1$ and $G_{kp} = 0 \forall k \neq j$ for any designated p . Hence, it is not necessary to impose such a restriction for every factor. This restriction would be used when the user believes that the j th outcome is the p th factor but measured with error. It may be fruitful to combine this restriction with the restriction that $\beta_{jq} = 0 \forall q \neq p$ but doing so is not required. Thus, this restriction is weaker and presumably more realistic than the restriction that the j th variable is “exogenous” and measured without error, i.e. $\beta_{jp} = \mathbb{I}\{q = p\}$.

These indicators can be specified interactively in response to the following dialog:⁴

⁴This restriction can also be enforced from the command line with `make_restrictions(*, criteria = list("indicators_1st"), methodArgs = list(indicators_1st = z))`, where z is an integer vector with r elements, each of which contains the row number of the best indicator of the p th factor, which can be `NA_integer_` to avoid constraining the p th factor.



The function that imposes this constraint is formally defined as the average column-wise rank of the designated indicators (among factors with designated indicators) less two. This function ranges from -1 to $n - 2$ and equals -1 if and only if the indicators have the largest factor contribution in their respective columns.

- “Reference factors at level 1 must have more effective variance than does the battery as a whole”. The function that imposes this constraint is formally defined as⁵

$$\begin{cases} -1 & \text{if } |\Psi|^{\frac{1}{r}} \geq |\mathbf{R}|^{\frac{1}{n}} \\ |\mathbf{R}|^{\frac{1}{n}} - |\Psi|^{\frac{1}{r}} & \text{otherwise} \end{cases} \in [-1, 1],$$

This criterion loosely implies that the reference factors at level 1 be “less correlated” than is the configuration of outcomes in common factor space and encourages the primary axes to cut through the edges of the test configuration. This restriction is extremely weak and could be imposed in a more “exploratory” SEFA.

- “Primary factors at level 1 must have more effective variance than does the battery as a whole”. The function that imposes this constraint is formally defined as⁶

$$\begin{cases} -1 & \text{if } |\Phi|^{\frac{1}{r}} \geq |\mathbf{R}|^{\frac{1}{n}} \\ |\mathbf{R}|^{\frac{1}{n}} - |\Phi|^{\frac{1}{r}} & \text{otherwise} \end{cases} \in [-1, 1],$$

and is similar to the previous criterion above except is more intuitive because the effective variance of the primary factors is at issue. In general, it seems that imposing this restriction on the primary factors is *stronger* than imposing it on the reference factors and tends to push the first-order axes farther out into the edges of the test configuration. It seems clear that Yates (1987) would have supported this restriction but does not specifically mention it.

- “Communalities must exceed all the factor contributions that constitute them.” The function that imposes this constraint is formally defined as⁷

$$\frac{-1}{nr} \sum_{j=1}^n \sum_{p=1}^r \mathbb{I}\{h_j^2 \geq \mathfrak{D}_{jp}\} \in [-1, 0],$$

which permits some negative suppressor variables but prohibits extreme situations such as that where one factor contribution is large and positive and another in that row is almost equally large in magnitude but negative. Overall, this restriction seems quite weak.

⁵This constraint can be enforced by from the command line with `make_restrictions(*, criteria = list("ev_RF_1st"))`.

⁶This constraint can be enforced by from the command line with `make_restrictions(*, criteria = list("ev_PF_1st"))`.


⁷This constraint can be enforced by from the command line with `make_restrictions(*, criteria = list("h2_over_FC_1st"))`.

- “Best indicator of each factor must have non-negative factor contributions.” Let \mathcal{D} be the $r \times r$ submatrix of \mathcal{D} whose rows contain the best indicators of each factor as defined by the column-maximums of \mathcal{D} with the additional proviso that the rows are unique. The function that imposes this constraint is formally defined as⁸

$$\frac{-1}{r^2} \sum_{q=1}^r \sum_{p=1}^r \mathbb{I} \left\{ \mathcal{D}_{qp} \geq 0 \right\} \in [-1, 0],$$

and is stronger than the previous criterion and weaker than the next criterion. Although it prohibits negative suppressor variables among the best indicators, it permits any degree of suppression among the other outcome variables.

- “No negative suppressor variables.” Let $\mathcal{D}^* \leq 0$ be the minimum factor contribution that the user is willing to accept, and thus wants all factor contributions to exceed this threshold to prohibit “negative” suppressor variables but perhaps make some allowance for factor contributions to be slightly negative by chance. This constraint can be specified interactively via the following dialog with $\mathcal{D}^* = -0.01$.⁹



The function that imposes this constraint is formally defined as

$$\frac{-1}{nr} \sum_{j=1}^n \sum_{p=1}^r \mathbb{I} \{ \mathcal{D}_{jp} \geq \mathcal{D}^* \} \in [-1, 0],$$

and is a fairly strong — yet in many cases, reasonable — restriction. Yates (1987, p. 119) essentially recommends it, albeit with some incorrect statements about the relationship between this restriction and the rest of the surrounding chapter.

- “Reference factors must have more generalized variance than do primary factors”. The function that imposes this constraint could be formally defined as¹⁰

$$\begin{cases} -1 & \text{if } |\Phi| \leq |\Psi| \\ |\Phi| - |\Psi| & \text{otherwise} \end{cases} \in [-1, 1]$$

but has a faster and substantively equivalent operationalization in the code. When $r_1 = 2$, $|\Phi| = |\Psi|$ by necessity, so this criterion can bind in the case when $r \geq 3$. Also, this criterion necessarily holds when there is a single second-order factor. Yates (1987, p. 27) asserts that this restriction is a necessary condition of a useful factor analysis model, although counter-examples can be generated. With randomly drawn correlations from a uniform distribution on $[-1, 1]$, Φ is both positive definite and satisfies this restriction about 25% of the time when $r = 3$, 5% of the time when $r = 4$, 0.5% of the time when $r = 5$, and even less frequently for $r > 5$.

⁸This constraint can be enforced from the command line with `make_restrictions(*, criteria = list("distinguishability_1st"))`.

⁹This constraint can be enforced at the command line with `make_restrictions(*, criteria = list("no_neg_suppressors_1st"), methodArgs = list(FC_threshold_1st = -.01))`


¹⁰This constraint can be enforced by from the command line with `make_restrictions(*, criteria = list("gv_1st"))`.

- “Outcomes in hyperplanes have more effective variance than does the battery as a whole”. The function that imposes this constraint is formally defined as¹¹

$$\frac{-1}{r} \sum_{p=1}^r \mathbb{I} \left\{ \left| \beta^p \Phi \beta'^p + \Theta^2 \right|^{\frac{1}{b_p}} \geq |\mathbf{R}|^{\frac{1}{n}} \right\} \in [-1, 0],$$

where β^p is the $b_p \times r$ submatrix of β with exact zeros in the p th column and Θ^2 is the $b_p \times b_p$ diagonal submatrix of Θ^2 with the unique variances for these b_p outcomes. Thus, this criterion loosely requires that the outcomes in the p th hyperplane be “less correlated” than is the battery as a whole (in common factor space) for all r hyperplanes and is thus fairly strong. This constraint goes well with the revisionist Yates mapping rule discussed in section 6.2.5.

- “Minimum binary distance among columns of the logically negated loadings matrix.” Let $\beta_{jp}^* = \begin{cases} 1 & \text{if } \beta_{jp} = 0 \\ 0 & \text{if } \beta_{jp} \neq 0 \end{cases}$ be the “logically negated loadings matrix”. The binary distance between two columns of the “logically negated loadings matrix” is defined as (see ?dist) as “the *proportion* of bits in which only one is on amongst those in which at least one is on”. In order to satisfy the theorem in Reiersøl (1950), the minimum binary distance between any two columns must be positive, but it is possible to require a larger minimum binary distance. Let d^* be the requisite minimum binary distance, which can be specified in response to the following dialog:



Let d_{qp} be the binary distance between the q th and p th columns of β . The function that imposes this constraint is formally defined as¹²

$$\frac{-2}{r(r-1)} \sum_{q=1}^r \sum_{p \neq q}^r \mathbb{I} \{d_{qp} \geq d^*\} \in [-1, 0]$$

and would most often be used in a more “exploratory” SEFA.

- “Generalized variance of the reproduced correlation matrix must be less than that of the sample correlation matrix.” The function that imposes this constraint is formally defined as¹³

$$\begin{cases} -1 & \text{if } |\mathbf{R}| \geq |\text{cor}(\mathbf{Y})| \\ |\text{cor}(\mathbf{Y})| - |\mathbf{R}| & \text{otherwise} \end{cases} \in [-1, 1],$$

where $|\text{cor}(\mathbf{Y})|$ is the determinant of the sample correlation matrix. This restriction is pretty weak.

- “Prohibit some coefficients from being zero.” This restriction is relevant only for SEFA, where it is possible that the mapping rule will map some coefficient to zero that theoretically should not be zero. The most common case where this restriction is helpful is when equality restrictions are impose on two coefficients, and one presumably intends that they both be nonzero. The genetic algorithm would otherwise tend to make both zero and thereby satisfy two restrictions the easy way.

¹¹This constraint can be enforced from the command line with `make_restrictions(*, criteria = list("cohyperplanarity_1st"))`.

¹²This constraint can also be enforced from the command line with `make_restrictions(*, critieria = list("dist_cols_1st"), methodArgs = list(cutpoint_1st = 0.5))`

¹³This constraint can be enforced from the command line with `make_restrictions(*, criteria = list("volume_1st"))`.

Let β^* be a matrix such that $\beta_{jp}^* = 1$ if β_{jp} must not be zero and equals zero otherwise, which can be specified in response to the following dialog:

Row.names	F1	F2	F3	F4	F5
VisualPerception	FALSE	FALSE	FALSE	FALSE	FALSE
Cubes	FALSE	FALSE	FALSE	FALSE	FALSE

The function that imposes this constraint is formally defined as¹⁴

$$\frac{-1}{\sum_{j=1}^n \sum_{p=1}^r \beta_{jp}^*} \sum_{j=1}^n \sum_{p=1}^r \mathbb{I}\{\beta_{jp}^* = 1\} \mathbb{I}\{\beta_{jp} \neq 0\} \in [-1, 0]$$

and equals -1 if and only if none of the prohibited coefficients are zero.

Note that it is possible to select (or specify at the command line) more than one of these criteria as constraints in the lexical optimization process.

¹⁴This constraint can also be enforced at the command line with `make_restrictions(*, criteria = list("block_1st"), methodArgs = list(blockers = beta_star))`

7 Exploratory Factor Analysis via `Factnal()` and `Rotate()`

To start this sequence, I typed the command

```
man <- make_manifest(covmat = Harman74.cor)
res <- make_restrictions(manifest = man, factors = 5, model = "EFA")
EFA <- Factnal(manifest = man, restrictions = res)
```

7.1 Factor Extraction

I use Λ to indicate the preliminary pattern matrix to estimate when the preliminary factors are orthogonal. The above sequence of commands will estimate the EFA model under the assumptions that $\Phi = \mathbf{I}$ and that $\Lambda' \Theta^{-2} \Lambda$ is diagonal using the maximum likelihood discrepancy function. If a different discrepancy function were specified, then the EFA model would be estimated under the assumptions that $\Phi = \mathbf{I}$ and that $\Lambda = \begin{bmatrix} \Lambda_1 \\ \Lambda_2 \end{bmatrix}$ where Λ_1 is a $r \times r$ matrix with zeros above its diagonal.

7.2 Factor Transformation

Note that regardless of how the EFA model was estimated, the preliminary factors will have the noncanonical form where $\Lambda = \begin{bmatrix} \Lambda_1 \\ \Lambda_2 \end{bmatrix}$ and Λ_1 is a $r \times r$ matrix with zeros above its diagonal. After verifying that a EFA model with r factors fits the data reasonably well, there were no near-Heywood cases, etc., the next task is to choose a transformation of the preliminary factors.

Let $\Phi = \mathbf{T}'\mathbf{T}$ where \mathbf{T} is a $r \times r$ transformation matrix with unit-length columns (not rows). The factor transformation step chooses \mathbf{T} so so that some (lexical) objective function is *minimized*. I initiated this sequence with

```
EFA_rotated <- Rotate(EFA)
```

WARNING: The subsequent pop-up menus are presented here in the logical order, which is different than the order in which they will appear when calling `Rotate()`.

7.2.1 Functional inequality restrictions


Again let $\mathbb{I}\{\cdot\}$ be the “indicator function”, which equals 1 if the condition within the braces is true and equals 0 if the condition is false. In FAiR, an inequality restriction function is a piecewise function that equals -1 if the restriction is satisfied and equals some number greater than -1 if the restriction is not satisfied. You do not need to worry too much about the return values for the case where the restriction is not satisfied.

One such inequality restriction is mandatory and is used to prevent factor collapse. The function that imposes this constraint is formally defined as

$$\begin{cases} -1 & \text{if } |\Phi|^{\frac{1}{r}} \geq v^* \\ v^* - |\Phi|^{\frac{1}{r}} & \text{otherwise} \end{cases} \in [-1, 1],$$


where $v^* > 0$ is the minimum acceptable effective variance for the primary factors. Please refer to section 4 for the definition of effective variance. One can specify v^* by responding to the following dialog:¹⁵

¹⁵The corresponding pop-up menu can be avoided from the command line with `Rotate(EFA, criteria = list("no_factor_collapse"), methodArgs = list(nfc_threshold = .3))`




A value of $v^* = 0.3$ is quite weak and in the $r = 2$ case would permit the off-diagonal of Φ to be up to ± 0.954 , which only precludes *literal* factor collapse. One deliberately nice property of defining the restriction with respect to the effective variance is that the same v^* is more-or-less reasonable regardless of r .

The following menu is similar to that in section 6.2.7, which offered choices for functional inequality restrictions on SEFA and CFA models. Except for the first two choices, the definitions of the criteria and subsequent pop-up menus are the same; only the two that differ will be discussed in this subsection.




- “Bounds on correlations among primary factors.” This restriction is conceptually simple and merely places lower and / or upper bounds on all off-diagonals of $\Phi = \mathbf{T}'\mathbf{T}$. However, because the objective function is minimized over the cells of \mathbf{T} instead of Φ , we have to use a constraint function to accomplish this goal. Let $\underline{\Phi}$ be the minimum acceptable factor correlation and let $\bar{\Phi}$ be the maximum acceptable factor correlation, which can be specified interactively as follows:¹⁶

¹⁶This constraint can also be enforced from the command line with `Rotate(EFA, criteria = list("limit_correlations"), methodArgs(lower = -.1, upper = 1))`



The function that imposes this constraint returns -1 if $\underline{\Phi} \leq \Phi_{pq} \leq \bar{\Phi} \forall p > q$ and otherwise returns the distance between the relevant bound and the correlation that most egregiously fails to satisfy these bounds.

- “Positive manifold.” Similarly, this restriction merely places a lower bound on $\Upsilon_{jp} \forall j, p$. Let $\Upsilon^* \leq 0$ be this lower bound, which can be specified interactively by responding to the following dialog:¹⁷




The function that imposes this constraint is formally defined as

$$\frac{-1}{nr} \sum_{j=1}^n \sum_{p=1}^r \mathbb{I}\{\Upsilon_{jp} \geq \Upsilon^*\} \in [-1, 0]$$

and hence is satisfied if and only if the proposed \mathbf{T} yields a “positive” manifold.

7.2.2 The ultimate analytic criterion




The next step is to choose the ultimate criterion for lexical optimization, as shown in the above dialog box. These criteria are defined as follows:

- phi: This criterion was proposed by Thurstone (1935) to numerically characterize simple structure when it reaches its theoretical minimum of (log) zero. The criterion is formally defined as


$$\ln \left(\sum_{j=1}^n \exp \left(\frac{1}{c} \sum_{p=1}^{r_1} \ln (\Upsilon_{jp}^2) \right) \right),$$

¹⁷This constraint can be enforced at the command line with `Rotate(EFA, criteria = list("positive_manifold"), methodArgs(pm_threshold = -0.1))`

which is equal to logarithm of ϕ as defined by Thurstone (1935) when $c = 1$. Thurstone suggested that choosing c to be greater than 1.0 could yield better results by lessening the pressure to get one exact zero into each row of Υ and increasing the pressure to get more than one near-zero into some rows of Υ . If Thurstone's criterion is used, the following dialog box will appear, prompting the user for c :¹⁸



- **varphi**: This criterion is a generalization of Thurstone's ϕ . Let $\vec{\Upsilon}$ be a matrix of reference structure correlations where each row is independently sorted in increasing magnitude, and let $\vec{\phi}_{-[1:p]}$ be Thurstone's criterion (with $c = 1$) calculated on $\vec{\Upsilon}$, excluding the first through the p th column of $\vec{\Upsilon}$. Then define $\varphi = \ln \left(\phi + \sum_{p=1}^{r-2} w_p \vec{\phi}_{-[1:p]} \right)$, where $w_p \in [0, 1]$ is the weight the analyst specifies for $\vec{\phi}_{-[1:p]}$ relative to a unit weight placed on ϕ . At one extreme, if $w_p = 0 \forall p$, then $\varphi = \phi$. At the other extreme, if $w_p = 1 \forall p$, then the analyst favors a perfect cluster configuration, where there is only one non-zero coefficient per test. An "objective" alternative to specifying weights is to specify that the weights are a function of $\vec{\Upsilon}$, such as $w_p = 1 - \max \left\{ \left(\vec{\Upsilon}_{p+1} \right)^2 \right\}^{\frac{1}{2}}$ to gradually reduce the weight as p increases. These weights can be specified interactively¹⁹:



- **minimaximin**: This criterion attempts to satisfy Thurstone's definition of simple structure by choosing \mathbf{T} so that all outcomes have at least one near-zero reference structure correlation. The criterion is formally defined as


$$\ln \left(\max \left\{ \min \left\{ \Upsilon_1^2 \right\}, \min \left\{ \Upsilon_2^2 \right\}, \dots, \min \left\{ \Upsilon_n^2 \right\} \right\} \right).$$

Thus, the (log of the) maximum, minimum, squared, row-wise reference structure correlation is minimized.

¹⁸This pop-up menu can be avoided from the command line with `Rotate(EFA, criteria = list("phi"), methodArgs(c = 1))`


¹⁹These pop-up menus can be avoided from the command line with `Rotate(EFA, criteria = list("varphi"), methodArgs(w = rep(0.5, 4)))` to weight all subsequent columns at half the weight of the first, for example. Or, `w` can be any negative number to indicate dynamic weights.

- LS: This criterion is called the Loading Simplicity Index in Lorenzo-Seva (2003). The derivation of it is not complicated but involves a lot of notation not previously introduced here, so the reader is referred to Lorenzo-Seva (2003) for details. Simply put, it reaches its optimum of (negative) 1.0 when Υ exhibits a perfect cluster configuration. Although Lorenzo-Seva (2003) defines the Loading Simplicity Index in terms of the primary pattern matrix, it is invariant to the column-scale of the primary pattern matrix and thus can arbitrarily be defined for the reference structure matrix as well.
- “Some oblique criterion from the GPArotation package.” If this option is marked, then the following dialog box appears, unless the name of the criterion is specified in `criteria` in the call to `Rotate`, e.g. `Rotate(EFA, criteria = list("quartimin"))`



These criteria are all defined in Browne (2001) and Bernaards and Jennrich (2006) and documented in the suggested GPArotation package. The versions in FAiR work the same way, with four exceptions. First, and most importantly, FAiR minimizes these criteria subjects to the aforementioned restrictions. Second, if the criterion has any additional arguments, they can be specified in a corresponding pop-up menu (or specified in `methodArgs`, as in the GPArotation library). Third, the partially-specified target implementation allows the partially-specified target matrix to have NA in cells that are not targeted, instead of specifying a separate weight matrix of zeros and ones. Fourth, in FAiR the criteria defined in the GPArotation package can also be invoked on

Υ or \mathcal{D} , instead of β by specifying `matrix = "RS"` or `matrix = "FC"` as an element of `methodArgs` or by responding to this dialog:



8 References

- Anderson, T.W. and H. Rubin. 1956. "Statistical inference in factor analysis" in *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability*. vol. 5. no. 1. University of California Press.
- Bentler, P.M. and P. Dudgeon. 1996. "COVARIANCE STRUCTURE ANALYSIS: Statistical Practice, Theory, and Directions." *Annual Reviews in Psychology*. 47(1):563 – 592.
- Bernaards, C.A. and R.I. Jennrich. 2006. "Gradient Projection Algorithms and Software for Arbitrary Rotation Criteria in Factor Analysis." *Educational and Psychological Measurement*. 65(5):676 – 696.
- Bollen, K.A. and K.G. Jöreskog. 1985. "Uniqueness does not Imply Identification: A Note on Confirmatory Factor Analysis." *Sociological Methods and Research*. 14(2):155 – 163.
- Browne, M.W. 2001. "An Overview of Analytic Rotation in Exploratory Factor Analysis." *Multivariate Behavioral Research*. 36(1):111 – 150.
- Butler, J.M. 1969. "Simple Structure Reconsidered: Distinguishability and Invariance in Factor Analysis." *Multivariate Behavioral Research* 4(1):5 – 28.
- Conger, J.A. 1974. "A Revised Definition for Suppressor Variables: a Guide To Their Identification and Interpretation." *Educational and Psychological Measurement*. 34(1):35 – 46
- Dey, D. K. and K. Srinivasan. 1985. "Estimation of a covariance matrix under Stein's loss." *The Annals of Statistics*. 13:1581 – 1591.
- Goodrich, B. 2008a. "SEFAiR So Far." Unpublished manuscript available from the FAiR wiki <http://wiki.r-project.org/rwiki/doku.php?id=packages:cran:fair>
- Goodrich, B. 2008b. "Restrictions, Factor Analysis, and Genetic Algorithms." Unpublished manuscript available from the FAiR wiki <http://wiki.r-project.org/rwiki/doku.php?id=packages:cran:fair>
- Harman, H.H. 1976. *Modern Factor Analysis*. University Of Chicago Press.
- Howe, W.G. 1955. Some Contributions to Factor Analysis. Technical report ORNL-1919, Oak Ridge National Lab., Tenn.
- Koopmans, T.C. and O. Reiersøl. 1950. "The identification of structural characteristics." *Annals of Mathematical Statistics*. 21(1):165 – 181.
- Lorenzo-Seva, U. 2003. "A factor simplicity index." *Psychometrika* 68(1):49 – 60.

- Millsap, R.E. 2001, "When Trivial Constraints Are Not Trivial: The Choice of Uniqueness Constraints in Confirmatory Factor Analysis." *Structural Equation Modeling*. 8(1):1 – 17.
- Opgen-Rhein, R., and K. Strimmer. 2007. "Accurate ranking of differentially expressed genes by a distribution-free shrinkage approach." *Statistical Applications in Genetics and Molecular Biology*. 6(1), article 9.
- Peña, D. and J. Rodríguez. 2003. "Descriptive measures of multivariate scatter and linear dependence." *Journal of Multivariate Analysis* 85(2):361 – 374.
- Pison, G., Van Aelst, S., and Willems, G. 2002. "Small Sample Corrections for LTS and MCD." *Metrika*. 55:111 – 123.
- Reiersøl, O. 1950. "On the Identifiability of Parameters in Thurstone's Multiple Factor Analysis." *Psychometrika*. 15:121 – 149.
- Rousseeuw, P.J. and K. Van Driessen. 1999. "A fast algorithm for the minimum covariance determinant estimator." *Technometrics*. 41(3):212 – 223.
- Schöfer, J., and K. Strimmer. 2005. "A shrinkage approach to large-scale covariance matrix estimation and implications for functional genomics." *Statistical Applications in Genetics and Molecular Biology*. 4(1), article 32.
- Thurstone, L.L. 1935. *The vectors of mind: multiple-factor analysis for the isolation of primary traits*. University of Chicago Press.
- White, O. 1966. "Some Properties Of Three Factor Contribution Matrices." *Multivariate Behavioral Research*. 1(3):373 – 377.
- Yates, A. 1987. *Multivariate exploratory data analysis: A perspective on exploratory factor analysis*. State University of New York Press.